

Alexa Fluor® 647 anti-STAT3 Phospho (Ser727) Antibody

Catalog# / Size	698913 / 25 tests 698914 / 100 tests
Clone	A16089B
Regulatory Status	RUO
Other Names	Signal transducer and activator of transcription 3, Acute-phase response factor (APRF), HIES, ADMIO
Isotype	Mouse IgG1, κ
Description	<p>The STAT3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors and is required for murine fetal development. STAT3 is an 88 kD member of the STAT (signal transducer and activators of transcription) protein family that is phosphorylated in response to a cytokine receptor-associated kinase activity. Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding. Transcriptional activation was reported to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways.</p>

STAT3 forms both homo- and heterotrimers and is involved in the activation of genes required for cell growth and apoptosis. STAT3 is also involved in gp130 signaling and binds to IL-6 response elements in various acute phase protein promoters. STAT3 is phosphorylated by signaling from IFNs, EGF, FGF, IL-5, HGF, LIF, and BMP2. STAT3 activity is inhibited by PIAS3 and GRIM-19 and can also be regulated by the Rac1 protein.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human Stat3 peptide phosphorylated at Ser 727. Complete Freund's adjuvant.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by intracellular flow cytometry using our True-Phos™ Perm Buffer in Cell Suspensions Protocol. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.</p> <p>* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.</p> <p>Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.</p> <p>View full statement regarding label licenses</p>
Excitation Laser	Red Laser (633 nm)
Application Notes	The STAT3 Phospho (Ser727) antibody recognizes the regulatory serine phosphorylation of human STAT3 protein.
Product Citations	1. Martínez-Fábregas J, <i>et al.</i> 2020. Cell Rep. 33:108545. PubMed

RRID

AB_2750259 (BioLegend Cat. No. 698913)
AB_2750260 (BioLegend Cat. No. 698914)

Antigen Details

Structure	STAT3 is a 770 amino acid protein of 88 kD. It consists of a DNA binding domain, a SH2 domain, a regulatory tyrosine responsible for binding of SH2 domain, and a C-terminal transactivation domain.
Distribution	Ubiquitous
Function	STAT3 is also involved in gp130 signaling and binds to IL-6 response elements in various acute phase protein promoters. STAT3 is phosphorylated by signaling from IFNs, EGF, FGF, IL-5, HGF, LIF, and BMP2. STAT3 activity is inhibited by PIAS3 and GRIM-19 and can also be regulated by the Rac1 protein.
Interaction	When activated, STAT3 dimerizes and translocates to the nucleus where it regulates gene expression.
Biology Area	Cell Biology, Neuroscience, Neuroscience Cell Markers, Signal Transduction, Transcription Factors
Molecular Family	Nuclear Markers, Phospho-Proteins
Antigen References	1. Akira S, <i>et al.</i> 1994. <i>Cell</i> . 77:63. 2. Zhang X, <i>et al.</i> 1995. <i>Science</i> 267:1990. 3. Sanchez-Margalet V, <i>et al.</i> 2001. <i>Cell. Immunol.</i> 211:30. 4. Simon A, <i>et al.</i> 2000. <i>Science</i> 290:144. 5. Hoey T, <i>et al.</i> 1999. <i>Adv. Immunol.</i> 71:145.
Gene ID	6774

Related Protocols


[Intracellular Staining With True-Phos™ Perm Buffer in Cell Suspensions Protocol](#)

[Intracellular Staining With True-Phos™ Perm Buffer in Whole Blood](#)

Other Formats

Go-ChIP-Grade™ Purified anti-STAT3 Phospho (Ser727), PE anti-STAT3 Phospho (Ser727), Purified anti-STAT3 Phospho (Ser727), PerCP/Cyanine5.5 anti-STAT3 Phospho (Ser727), PE/Cyanine7 anti-STAT3 Phospho (Ser727), Alexa Fluor® 488 anti-STAT3 Phospho (Ser727), Brilliant Violet 421™ anti-STAT3 Phospho (Ser727), Alexa Fluor® 647 anti-STAT3 Phospho (Ser727)

Product Data



Human peripheral blood monocytes were treated with (filled histogram), or without (open histogram), Cell Activation Cocktail (Cat. No. 423301) for 15minutes, fixed with Fixation Buffer (Cat. No. 420801), permeabilized with True-Phos™ Perm Buffer (Cat. No. 425401), then intracellularly stained with Human TruStain FcX™ (Cat. No. 422302), True-Stain Monocyte Blocker™ (Cat. No. 426103), anti-STAT3 Phospho (Ser727) (clone A16089B) Alexa Fluor® 647.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587