

APC/Fire™ 810 anti-mouse CD19 Antibody

Catalog# / Size	115577 / 25 µg 115578 / 100 µg
Clone	6D5
Regulatory Status	RUO
Other Names	B4
Isotype	Rat IgG2a, κ
Description	CD19 is a 95 kD glycoprotein also known as B4. It is a member of the Ig superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Mouse CD19-expressing K562 human erythroleukemia cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 810 under optimal conditions.
Concentration	0.2 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>* APC/Fire™ 810 has a maximum excitation of 650 nm and a maximum emission of 810 nm.</p> <p>Excessive exposure to light, and commonly used fixation, permeabilization buffers can affect APC/Fire™ 810 fluorescence signal intensity and spread. Please keep conjugates protected from light exposure. For more information and representative data, visit our Fire Dyes page.</p>
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunofluorescence ⁷ .
Application References	<ol style="list-style-type: none"> Shoham T, <i>et al.</i> 2003. <i>J. Immunol.</i> 171:4062. (FC) Goodyear CS, <i>et al.</i> 2004. <i>J. Immunol.</i> 172:2870. (FC) Kamimura D, <i>et al.</i> 2006. <i>J. Immunol.</i> 177:306. (FC) Andoniou CE, <i>et al.</i> 2005. <i>Nat. Immunol.</i> 6:1011. (FC) Lawson BR, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:5366. (FC) Phan TG, <i>et al.</i> 2007. <i>Nat. Immunol.</i> 8:992. (FC) Hayashida K, <i>et al.</i> 2008. <i>J. Biol. Chem.</i> 283:19895. (IF) PubMed Charles N, <i>et al.</i> 2010. <i>Nat. Med.</i> 16:701. (FC) PubMed Bankoti J, <i>et al.</i> 2010. <i>Toxicol. Sci.</i> 115:422. (FC) PubMed Stadnisky MD, <i>et al.</i> 2011. <i>Blood.</i> 117:5133. (FC) PubMed Perlot T, <i>et al.</i> 2012. <i>J. Immunol.</i> 188:1201. (FC) PubMed Olive V, <i>et al.</i> 2013. <i>Elife.</i> 2:822. PubMed
(PubMed link indicates BioLegend citation)	

RRID AB_2892274 (BioLegend Cat. No. 115577)
 AB_2892274 (BioLegend Cat. No. 115578)

Antigen Details

Structure	Ig superfamily, associates with CD21 and CD81, 95 kD
Distribution	Pro-B cells to mature B cells (during development), follicular dendritic cells
Function	Modulates B cell activation and differentiation
Ligand/Receptor	CD21, CD81, Leu-13
Cell Type	B cells, Dendritic cells
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	1. Fearon DT. 1993. <i>Curr. Opin. Immunol.</i> 5:341. 2. Krop I, et al. 1996. <i>Eur. J. Immunol.</i> 26:238. 3. Krop I, et al. 1996. <i>J. Immunol.</i> 157:48. 4. Tedder TF, et al. 1994. <i>Immunol. Today</i> 15:437.
Gene ID	12478


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse CD19, Biotin anti-mouse CD19, FITC anti-mouse CD19, PE anti-mouse CD19, PE/Cyanine5 anti-mouse CD19, Purified anti-mouse CD19, PE/Cyanine7 anti-mouse CD19, Alexa Fluor® 488 anti-mouse CD19, Alexa Fluor® 647 anti-mouse CD19, Pacific Blue™ anti-mouse CD19, Alexa Fluor® 700 anti-mouse CD19, APC/Cyanine7 anti-mouse CD19, PerCP anti-mouse CD19, PerCP/Cyanine5.5 anti-mouse CD19, Alexa Fluor® 594 anti-mouse CD19, Brilliant Violet 421™ anti-mouse CD19, Brilliant Violet 570™ anti-mouse CD19, Brilliant Violet 605™ anti-mouse CD19, Brilliant Violet 650™ anti-mouse CD19, Brilliant Violet 785™ anti-mouse CD19, Brilliant Violet 510™ anti-mouse CD19, Purified anti-mouse CD19 (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse CD19, Brilliant Violet 711™ anti-mouse CD19, APC/Fire™ 750 anti-mouse CD19, TotalSeq™-A0093 anti-mouse CD19, Brilliant Violet 750™ anti-mouse CD19, TotalSeq™-B0093 anti-mouse CD19, Spark Blue™ 550 anti-mouse CD19, Spark NIR™ 685 anti-mouse CD19, TotalSeq™-C0093 anti-mouse CD19, Ultra-LEAF™ Purified anti-mouse CD19, PE/Fire™ 640 anti-mouse CD19 Antibody, Spark YG™ 581 anti-mouse CD19, APC/Fire™ 810 anti-mouse CD19, Spark YG™ 570 anti-mouse CD19, Spark Blue™ 574 anti-mouse CD19 Antibody

Product Data



C57BL/6 mouse splenocytes were stained with anti-mouse CD3 FITC and anti-mouse CD19 (clone 6D5) APC/Fire™ 810 (left) or rat IgG2a, κ APC/Fire™ 810 isotype control (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587