

## Spark Red™ 718 anti-mouse CD274 (B7-H1, PD-L1) Antibody

<b>Catalog# / Size</b>	124347 / 25 µg 124348 / 100 µg
<b>Clone</b>	10F.9G2
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	B7-H1, PD-L1
<b>Isotype</b>	Rat IgG2b, κ
<b>Description</b>	CD274, also known as B7-H1 or programmed death ligand 1 (PD-L1), is a 40 kD type I transmembrane protein and a member of the B7 family within the immunoglobulin receptor superfamily. It is expressed on T cells, B cells, NK cells, dendritic cells, IFN-γ activated endothelial cells, and monocytes. B7-H1 is one of the ligands of PD-1. The interaction of B7-H1 with PD-1 plays an important role in the inhibition of T cell responses. Other studies have shown that B7-H1 is able to costimulate T cell growth and cytokine production. CD274 is involved in costimulation essential for T cell proliferation and production of IL-10 and IFN-γ, in an IL-2-dependent and a PD-1-independent manner. Its interaction with PD-1 inhibits T cell proliferation and cytokine production.

### Product Details

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with Spark Red™ 718 under optimal conditions.
<b>Concentration</b>	0.5 mg/mL
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.  * Spark Red™ 718 has a maximum excitation of 697 nm and a maximum emission of 718 nm.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunofluorescence <sup>4</sup> , blocking <sup>6,7,8,9</sup> , and immunohistochemistry of acetone-fixed frozen sections <sup>4, 11</sup> . The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 124303). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 124318) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).
<b>Application References</b>	<ol style="list-style-type: none"> <li>1. Maier H, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:2714.</li> <li>2. Meng Q, <i>et al.</i> 2006. <i>Invest. Ophthalmol. Vis. Sci.</i> 47:4444. <a href="#">PubMed</a></li> <li>3. Scarlett UK, <i>et al.</i> 2012. <i>J Exp Med.</i> 209:495. <a href="#">PubMed</a></li> <li>4. Grabie N, <i>et al.</i> 2007. <i>Circulation</i> 116:2062. (IF, IHC)</li> <li>5. Paterson AM, <i>et al.</i> 2011. <i>J. Immunol.</i> 187:1097.</li> <li>6. Channappanavar R, <i>et al.</i> 2012. <i>PLoS One</i> 7:e39757. (Block)</li> <li>7. Schreiber HA, <i>et al.</i> 2010. <i>PLoS One</i> 5:e11453. (Block) <a href="#">PubMed</a></li> <li>8. Muthumani K, <i>et al.</i> 2011. <i>J. Immunol.</i> 187:2932. (Block) <a href="#">PubMed</a></li> </ol>
<b>(PubMed link indicates BioLegend citation)</b>	

9. Cripps JG, *et al.* 2010. *Hepatology* 52:1350. (Block) [PubMed](#)
10. Murakami R, *et al.* 2013. *PLoS One.* 8:73270. [PubMed](#)
11. Riella LV, *et al.* 2011. *Am. J. Transplant* 11:832-40. (IHC)
12. Lei GS, *et al.* 2015. *Infect Immun.* 83:572. [PubMed](#)

**RRID** AB\_2924463 (BioLegend Cat. No. 124347)  
 AB\_2924463 (BioLegend Cat. No. 124348)

## Antigen Details

<b>Structure</b>	40 kD type I transmembrane protein member of B7 family within the immunoglobulin receptor superfamily
<b>Distribution</b>	T cells, B cells, NK cells, dendritic cells, IFN- $\gamma$ activated endothelial cells, and monocytes
<b>Ligand/Receptor</b>	PD-1 (PDCD1)
<b>Cell Type</b>	B cells, Dendritic cells, Endothelial cells, Monocytes, NK cells, T cells
<b>Biology Area</b>	Cancer Biomarkers, Costimulatory Molecules, Immunology
<b>Molecular Family</b>	Adhesion Molecules, CD Molecules, Immune Checkpoint Receptors
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Sharpe A, <i>et al.</i> 2007. <i>Nat. Immunol.</i> 8:239.</li> <li>2. Dong H, <i>et al.</i> 1999. <i>Nat. Med.</i> 5:1365.</li> <li>3. Freeman G, <i>et al.</i> 2000. <i>J. Exp. Med.</i> 192:1027.</li> </ol>
<b>Gene ID</b>	<a href="#">60533</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-mouse CD274 (B7-H1, PD-L1), Biotin anti-mouse CD274 (B7-H1, PD-L1), PE anti-mouse CD274 (B7-H1, PD-L1), Brilliant Violet 421™ anti-mouse CD274 (B7-H1, PD-L1), APC anti-mouse CD274 (B7-H1, PD-L1), PE/Cyanine7 anti-mouse CD274 (B7-H1, PD-L1), Ultra-LEAF™ Purified anti-mouse CD274 (B7-H1, PD-L1), Brilliant Violet 711™ anti-mouse CD274 (B7-H1, PD-L1), Brilliant Violet 605™ anti-mouse CD274 (B7-H1, PD-L1), PE/Dazzle™ 594 anti-mouse CD274 (B7-H1, PD-L1), GoInVivo™ Purified anti-mouse CD274 (B7-H1, PD-L1), Brilliant Violet 785™ anti-mouse CD274 (B7-H1, PD-L1), PerCP/Cyanine5.5 anti-mouse CD274 (B7-H1, PD-L1), Brilliant Violet 650™ anti-mouse CD274 (B7-H1, PD-L1), PE/Cyanine5 anti-mouse CD274 (B7-H1, PD-L1), PE/Fire™ 640 anti-mouse CD274 (B7-H1, PD-L1), Spark Red™ 718 anti-mouse CD274 (B7-H1, PD-L1)

## Product Data



C57/B6 mouse splenocytes were stained with anti-mouse CD274 (B7-H1, PD-L1) (clone 10F.9G2) Spark Red™ 718 (filled histogram), or unstained control (open histogram).

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587