

TotalSeq™-B0564 anti-mouse Folate Receptor β (FR- β) Antibody

Catalog# / Size	153309 / 10 μ g
Clone	10/FR2
Regulatory Status	RUO
Other Names	FR- β
Isotype	Rat IgG2a, κ
Barcode Sequence	CTCAGATGCCCTTTA
Description	Folate receptor beta (FR- β) is a member of a family of folate binding receptors that have diverse structural identities but mediate transport of folates into cells. FR- β is a GPI-anchored protein with high affinity for folic acid. It is postulated that this receptor function as folate scavengers when folate supply is low or rapid cell growth requires elevated uptake of folate for methylation reactions including DNA biosynthesis. Although this protein was originally thought to be specific to placenta, it can also exist in other tissues, and it may play a role in the transport of methotrexate in synovial macrophages in rheumatoid arthritis patients.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Mouse Folate receptor β transfected cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-B oligomer under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-B antibodies are compatible with 10x Genomics Single Cell Gene Expression Solutions.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is $\leq 1.0 \mu$g per million cells in 100 μL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Additional Product Notes	<p>TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. 10x Genomics Chromium System and Reagents) and sequencer (e.g. Illumina analyzers) are required. Please contact technical support for more information, or visit biolegend.com/totalseq.</p> <p>The barcode flanking sequences are GTGACTGGAGTTCAGACGTGTGCTCTCCGATCTNNNNNNNNNN (PCR handle), and NNNNNNNNGCTTTAAGGCCGGTCCTAGC*A*A (capture sequence). N represents either</p>

randomly selected A, C, G, or T, and * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

Application References

(PubMed link indicates BioLegend citation)

1. Nagai T, et al. 2009. *Cancer Immunol Immunother.* 1577-86.

RRID AB_2876511 (BioLegend Cat. No. 153309)

Antigen Details

Structure	30 kD, seven disulfide bonds
Distribution	Placenta, monocytes, tissue resident macrophages
Function	Transport of folate into cells
Ligand/Receptor	Folic Acid
Cell Type	Macrophages, Monocytes
Antigen References	<ol style="list-style-type: none">1. Elwood PC, 1989, <i>J Biol Chem</i>, 264:14893.2. Shen F, et al, 1994, <i>Biochemistry</i>, 8;33:1209.3. Yi YS, 2016, <i>Immune Netw</i>,16:337.
Gene ID	14276

Related Protocols

[TotalSeq™-B or -C with 10x Feature Barcoding Technology](#)

Other Formats

Purified anti-mouse Folate Receptor β (FR- β), APC anti-mouse Folate Receptor β (FR- β), PE anti-mouse Folate Receptor β (FR- β), TotalSeq™-A0564 anti-mouse Folate Receptor β (FR- β), TotalSeq™-B0564 anti-mouse Folate Receptor β (FR- β) Antibody

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587