

## TotalSeq™-D Human Heme Oncology Cocktail, V1.0

<b>Catalog# / Size</b>	399906 / 8 tests
<b>Regulatory Status</b>	RUO
<b>Description</b>	The TotalSeq™-D Human Heme Oncology Cocktail, V1.0 has been designed to react with 42 unique cell surface antigens, including principal lineage antigens, and includes 3 isotype control antibodies, to aid in the multiomic characterization of immune cells.

The lyophilized cocktail provides convenience when processing several samples at the same time or at different time points, reducing inconsistencies associated with pipetting and handling multiplex vials and samples. The antibodies in the cocktail are provided at optimized concentrations to provide a ready-to-use solution once the cocktail has been reconstituted. The TotalSeq™-D Human Heme Oncology Cocktail, V1.0 comes packaged in convenient single-use vials.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Formulation</b>	Lyophilized from PBS containing stabilizers. Please reconstitute each tube with cell staining buffer as indicated in the application notes when ready to use. Make sure the cake is completely dissolved prior to using the cocktail for staining cells. The lyophilized cocktail material can vary in appearance and appear as either a lyophilized cake or a powder.
<b>Preparation</b>	This reagent is a combination of TotalSeq™-D oligo conjugated clones at optimal concentrations for single-cell sequencing analysis. See additional product notes for a list of clones.
<b>Storage &amp; Handling</b>	The lyophilized antibody cocktail should be stored between 2°C and 8°C in powder form and in a sealed container with desiccant until ready to use. Once vial is opened, it is to be reconstituted immediately. After reconstitution cocktail should be used within 2 hours.
<b>Application</b>	<a href="#">PG - Quality tested</a>
<b>Recommended Usage</b>	This panel has been optimized on human PBMCs for use with the Mission Bio Tapestri system. Using lysed whole blood or additional TotalSeq™ antibodies may require further optimization. Please contact our technical service group for further information.

Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.

<b>Application Notes</b>	<ol style="list-style-type: none"> <li>1. Equilibrate lyophilized panel to room temperature for 5 min.</li> <li>2. Spin down at 10,000 x g for 30 seconds at room temperature.</li> <li>3. Resuspend the lyophilized cocktail in 60 µL of Cell Staining Buffer (Cat. No. 420201). Replace the cap and vortex for 10 seconds. Note: Excess volume added to aid in removal of potential protein aggregates.</li> <li>4. Incubate at room temperature for 5 minutes.</li> <li>5. Vortex again and spin down at 10,000 x g for 30 seconds at room temperature.</li> <li>6. Transfer the entire volume (60 µL) of reconstituted panel to a low protein binding Eppendorf tube (Fisher cat. no. 022431081) or similar tube.</li> <li>7. Centrifuge at 14,000 x g for 15 min at 4°C.</li> <li>8. Proceed with staining the cells following <a href="#">Mission Bio User Guide</a>, Stain Cells section.</li> </ol>
--------------------------	---

Note: Staining cells with this cocktail should occur after cells have been stained with Human TruStain FcX™ (Fc Receptor Blocking Solution).

<b>Additional Product Notes</b>	TotalSeq™-D reagents are designed to profile protein expression at a single cell level. The Mission Bio Tapestri platform and DNA sequencer (e.g. Illumina sequencers) are required. Please contact <a href="#">technical support</a> for more information, or visit <a href="https://www.biolegend.com/en-us/totalseq/single-cell-dna">https://www.biolegend.com/en-us/totalseq/single-cell-dna</a>
---------------------------------	--

The barcode flanking sequences are CGAGATGACTACGCTACTCATGG (PCR handle), and

GAGCCGATCTAGTATCTCAGT\*C\*G (capture sequence). \* indicates a phosphorothioated bond, to prevent nuclease degradation.

The full oligomer sequence for this product, with the specific barcode in brackets is CGAGATGACTACGCTACTCATGG [Barcode] GAGCCGATCTAGTATCTCAGT\*C\*G.

TotalSeq-D0090 mouse IgG1, κ isotype ctrl has been observed to have elevated non-specific binding on monocytes in some donors. Isotype controls are included in the Heme Oncology Panel for the purpose of identifying low quality cells which have been observed to have staining for multiple isotype controls. It is not recommended to use isotype controls for thresholding of antibodies of the same isotype.



Download the excel file for a full list of clones and barcodes.

## Product Citations

1. Bianchi A, *et al.* 2022. *Genome Biol.* 23:229. [PubMed](#)

## RRID

AB\_2890849 (BioLegend Cat. No. 399906)

## Antigen Details


### Distribution

Multiple cell types and cell states


### Gene ID

NA

## Product Data



Human PBMCs were stained with the TotalSeq™-D Heme Oncology Cocktail v1.0 and processed using the Mission Bio Tapestry DNA and Protein workflow. Protein count data were transformed and visualized in a UMAP projection overlaid with protein. Clusters were identified based on protein expression only.



PBMCs from two donor samples were mixed together, stained with TotalSeq™-D antibodies and run on Mission Bio's Tapestry platform. Heatmap visualization reveals the power to obtain genotype and phenotype from the same cells across thousands of cells.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)).

BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587