

TotalSeq™-D0389 anti-human CD38 Antibody

Catalog# / Size	303553 / 10 µg
Clone	HIT2
Regulatory Status	RUO
Workshop	III 155
Other Names	T10, ADP-ribosyl cyclase
Isotype	Mouse IgG1, κ
Barcode Sequence	TGTACCCGCTTGTGA
Description	CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

Product Details

Verified Reactivity	Human
Reported Reactivity	Chimpanzee, Horse, Cow
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-D oligomer under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-D antibodies are compatible with Mission Bio's Tapestry Single-Cell Sequencing Platform for simultaneous detection of DNA and Protein.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Application Notes	Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections ⁶ and spatial biology (IBEX) ^{10,11} .
Additional Product Notes	TotalSeq™-D reagents are designed to profile protein expression at single cell level. The Mission Bio Tapestry platform and sequencer (e.g. Illumina analyzers) are required. Please contact technical support for more information, or visit biolegend.com/totalseq/single-cell-dna

The barcode flanking sequences are CGAGATGACTACGCTACTCATGG (PCR handle), and GAGCCGATCTAGTATCTCAGT*C*G (capture sequence). * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Application Technical Notes](#).

Application References

(PubMed link indicates BioLegend citation)

1. Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
2. Dieu M. 1998. *J. Exp. Med.* 188:373.
3. Esser M, *et al.* 2001. *J. Virol.* 75:6173.
4. Jeannin P, *et al.* 1999. *J. Immunol.* 162:2044.
5. Kapsogeorgou EK, *et al.* 2001. *J. Immunol.* 166:3107.
6. van der Voort R, *et al.* 1997. *J. Exp. Med.* 185:2121. (IHC)
7. Bende RJ, *et al.* 2003. *Am. J. Pathol.* 162:105.
8. Lehner M, *et al.* 2008. *J. Leukoc. Biol.* 83:883. [PubMed](#)
9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
10. Radtke AJ, *et al.* 2020. *Proc Natl Acad Sci USA.* 117:33455-33465. (SB) [PubMed](#)
11. Radtke AJ, *et al.* 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

RRID AB_2892351 (BioLegend Cat. No. 303553)

Antigen Details

Structure	ADP-ribosyl cyclase, ectoenzyme, type II glycoprotein, 45 kD
Distribution	T cells, B cells, NK, myeloid, plasma, and dendritic cells
Function	Ecto-ADP-ribosyl cyclase, calcium signaling, cell activation
Ligand/Receptor	CD31, hyaluronic acid
Cell Type	B cells, Dendritic cells, NK cells, Plasma cells, T cells
Biology Area	Immunology
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	1. Ferrero E, <i>et al.</i> 1999. <i>J. Leukoc. Biol.</i> 65:151. 2. Lund F, <i>et al.</i> 1995. <i>Immunol. Today</i> 16:469.
Gene ID	952

Related Protocols

[TotalSeq™-D with Mission Bio Tapestri® Single-Cell DNA + Protein Protocol](#)

Other Formats

APC anti-human CD38, FITC anti-human CD38, PE anti-human CD38, PE/Cyanine5 anti-human CD38, Purified anti-human CD38, Alexa Fluor® 488 anti-human CD38, Alexa Fluor® 647 anti-human CD38, PE/Cyanine7 anti-human CD38, Biotin anti-human CD38, PerCP anti-human CD38, PerCP/Cyanine5.5 anti-human CD38, Alexa Fluor® 700 anti-human CD38, Brilliant Violet 421™ anti-human CD38, Brilliant Violet 711™ anti-human CD38, Brilliant Violet 785™ anti-human CD38, Brilliant Violet 605™ anti-human CD38, APC/Cyanine7 anti-human CD38, Purified anti-human CD38 (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD38, Brilliant Violet 510™ anti-human CD38, TotalSeq™-A0389 anti-human CD38, TotalSeq™-C0389 anti-human CD38, APC/Fire™ 750 anti-human CD38, TotalSeq™-B0389 anti-human CD38, APC/Fire™ 810 anti-human CD38, Spark NIR™ 685 anti-human CD38 Antibody, TotalSeq™-D0389 anti-human CD38, GMP PE anti-human CD38, GMP FITC anti-human CD38, Pacific Blue™ anti-human CD38

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587