

Alexa Fluor® 488 anti-mouse CD80 Antibody

Catalog# / Size	104715 / 25 µg 104716 / 100 µg
Clone	16-10A1
Regulatory Status	RUO
Other Names	B7-1, B7, Ly-53
Isotype	Armenian Hamster IgG
Description	CD80 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD80 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD80 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD80 appears to be expressed later in the immune response than CD86. CD80 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

Product Details

Verified Reactivity	Mouse
Reported Reactivity	Dog
Antibody Type	Monoclonal
Host Species	Armenian Hamster
Immunogen	CHO cell line transfected with mouse B7 (CD80)
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 488 under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.</p> <p>Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.</p> <p>View full statement regarding label licenses</p>
Excitation Laser	Blue Laser (488 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunoprecipitation ² , <i>in vitro</i> and <i>in vivo</i> blocking of CTLA-4 Ig to CD80 by blocking costimulation of T cells by activated B cells ²⁻⁴ , and immunohistochemical staining of acetone-fixed frozen sections ^{1,4} . The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. Nos. 104747-104752).
Application References	1. Harlan DM, <i>et al.</i> 1994. <i>P. Natl. Acad. Sci. USA</i> 91:3137. (IHC)
(PubMed link indicates	2. Razi-Wolf Z, <i>et al.</i> 1992. <i>P. Natl. Acad. Sci. USA</i> 89:4210. (Block, IP)

- BioLegend citation)**
3. Hathcock KS, *et al.* 1994. *J. Exp. Med.* 180:631. (Block)
 4. Herold KC, *et al.* 1997. *J. Immunol.* 158:984. (Block, IHC)
 5. Ma XT, *et al.* 2006. *Cancer Res.* 66:1169.
 6. Andoniou CE, *et al.* 2005. *Nature Immunology* 6:1011. (FC)
 7. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366.
 8. Turnquist HR, *et al.* 2007. *J. Immunol.* 178:7018.
 9. Misra RS, *et al.* 2010. *J. Exp Med.* 207:1775. [PubMed](#)
 10. del Rio ML, *et al.* 2011. *Transpl. Int.* 24:501. (FC) [PubMed](#)
 11. Philipsen L, *et al.* 2013. *Mol Cell Proteomics.* 12:2551. [PubMed](#)

- Product Citations**
1. Espinosa-Cueto P, *et al.* 2017. *PLoS One.* 10.1371/journal.pone.0182126. [PubMed](#)
 2. Philipsen L, *et al.* 2013. *Mol Cell Proteomics.* 12:2551. [PubMed](#)
 3. Chanana P, *et al.* 2022. *Cells.* 11: [PubMed](#)
 4. Marangoni F, *et al.* 2021. *Cell.* [PubMed](#)
 5. Wong R, *et al.* 2020. *Immunity.* 53(5):1078-1094.e7. [PubMed](#)
 6. Liang J, *et al.* 2020. *Sci Adv.* 6:eabc3646. [PubMed](#)
 7. Yang BH, *et al.* 2020. *Cell Reports.* 27(12):3629-3645.e6. [PubMed](#)
 8. Chmielewski M and Abken H 2017. *Cell Rep.* 10.1016/j.celrep.2017.11.063. [PubMed](#)
 9. Wang L, *et al.* 2020. *J Nanobiotechnology.* 18:38. [PubMed](#)

RRID AB_492823 (BioLegend Cat. No. 104715)
AB_492822 (BioLegend Cat. No. 104716)

Antigen Details

Structure	Ig superfamily, 60 kD
Distribution	Macrophages, activated B cells and T cells, dendritic cells
Function	T cell costimulation
Ligand/Receptor	CD28 (stimulatory), CD152(CTLA4) (inhibitory)
Cell Type	B cells, Dendritic cells, Macrophages, T cells, Tregs
Biology Area	Cell Biology, Costimulatory Molecules, Immunology, Neuroscience, Neuroscience Cell Markers
Molecular Family	CD Molecules, Immune Checkpoint Receptors
Antigen References	<ol style="list-style-type: none">1. Barclay AN, <i>et al.</i> 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press.2. Linsley PS, <i>et al.</i> 1991. <i>J. Exp. Med.</i> 174:561.3. Salomon B, <i>et al.</i> 2001. <i>Annu. Rev. Immunol.</i> 19:225.
Gene ID	12519

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Biotin anti-mouse CD80, FITC anti-mouse CD80, PE anti-mouse CD80, Purified anti-mouse CD80, PE/Cyanine5 anti-mouse CD80, APC anti-mouse CD80, Alexa Fluor® 488 anti-mouse CD80, Alexa Fluor® 647 anti-mouse CD80, PerCP/Cyanine5.5 anti-mouse CD80, Pacific Blue™ anti-mouse CD80, Brilliant Violet 421™ anti-mouse CD80, Brilliant Violet 605™ anti-mouse CD80, Brilliant Violet 650™ anti-mouse CD80, PE/Cyanine7 anti-mouse CD80, Purified anti-mouse CD80 (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse CD80, APC/Fire™ 750 anti-mouse CD80, Brilliant Violet 711™ anti-mouse CD80, Brilliant Violet 510™ anti-mouse CD80, TotalSeq™-A0849 anti-mouse CD80, TotalSeq™-C0849 anti-mouse CD80, Ultra-LEAF™ Purified anti-mouse CD80, Alexa Fluor® 594 anti-mouse CD80, TotalSeq™-B0849 anti-mouse CD80 Antibody, PE/Fire™ 640 anti-mouse CD80, Spark NIR™ 685 anti-mouse CD80

Product Data

Thioglycolate-elicited Balb/c mouse
peritoneal macrophages stained with 16-
10A1 Alexa Fluor® 488



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587