

Biotin anti-mouse CD64 (FcγRI) Antibody

Catalog# / Size	139318 / 100 µg
Clone	X54-5/7.1
Regulatory Status	RUO
Other Names	FcRI
Isotype	Mouse IgG1, κ
Description	CD64 is a 72 kD single chain type I glycoprotein also known as FcγRI and FcRI. CD64 is a member of the immunoglobulin superfamily and is expressed on monocytes/macrophages, dendritic cells, and mast cells. The expression can be upregulated by IFN-γ stimulation. CD64 binds IgG immune complex. It plays a role in antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent cellular cytotoxicity (ADCC).

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	BALB/c mouse FcγRI-human IgG Fc fusion protein.
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤0.125 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	The X54-5/7.1 antibody reacts with mouse strains carrying CD64a and b alleles but not CD64d. X54-5/7.1 recognizes a conformational determinant formed between domains 2 and 3. Additional reported application (for relevant formats) include: immunoprecipitation ¹ , and spatial biology (IBEX) ^{5,6} . Clone X54-5/7.1 is not found to be useful for Western blots ¹ .
Application References	<ol style="list-style-type: none"> 1. Tan PS, <i>et al.</i> 2003. <i>J. Immunol.</i> 170:2549. (IP) 2. Ingersoll MA, <i>et al.</i> 2010. <i>Blood</i> 115:e10. (FC) 3. Ozeri E, <i>et al.</i> 2012. <i>J. Immunol.</i> 189:146. PubMed 4. Richardson ML, <i>et al.</i> 2014. <i>PLoS Negl Trop Dis.</i> 8:2825. PubMed 5. Radtke AJ, <i>et al.</i> 2020. <i>Proc Natl Acad Sci U S A.</i> 117:33455-65. (SB) PubMed 6. Radtke AJ, <i>et al.</i> 2022. <i>Nat Protoc.</i> 17:378-401. (SB) PubMed
Product Citations	<ol style="list-style-type: none"> 1. Shenoy A, <i>et al.</i> 2017. <i>PLoS Pathog.</i> 10.1371/journal.ppat.1006582. PubMed 2. Silva HM, <i>et al.</i> 2019. <i>J Exp Med.</i> 216:786. PubMed 3. Taniguchi A, <i>et al.</i> 2015. <i>Am J Physiol Lung Cell Mol Physiol.</i> 309: L789 - L800. PubMed
RRID	AB_2566557 (BioLegend Cat. No. 139318)

Antigen Details

Structure	Ig superfamily, type I glycoprotein, 72 kD
Distribution	Monocytes, macrophages, mast cells, dendritic cells
Function	Phagocytosis, ADCC
Ligand/Receptor	IgG
Cell Type	Dendritic cells, Macrophages, Mast cells, Monocytes
Biology Area	Immunology, Innate Immunity
Molecular Family	CD Molecules, Fc Receptors
Gene ID	14129



Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-mouse CD64 (FcγRI), PE anti-mouse CD64 (FcγRI), APC anti-mouse CD64 (FcγRI), PerCP/Cyanine5.5 anti-mouse CD64 (FcγRI), Brilliant Violet 421™ anti-mouse CD64 (FcγRI), Brilliant Violet 711™ anti-mouse CD64 (FcγRI), PE/Cyanine7 anti-mouse CD64 (FcγRI), FITC anti-mouse CD64 (FcγRI), Biotin anti-mouse CD64 (FcγRI), PE/Dazzle™ 594 anti-mouse CD64 (FcγRI), Alexa Fluor® 647 anti-mouse CD64 (FcγRI), Brilliant Violet 605™ anti-mouse CD64 (FcγRI), TotalSeq™-A0202 anti-mouse CD64 (FcγRI), TotalSeq™-C0202 anti-mouse CD64 (FcγRI), TotalSeq™-B0202 anti-mouse CD64 (FcγRI), PE/Cyanine5 anti-mouse CD64 (FcγRI), APC/Fire™ 750 anti-mouse CD64 (FcγRI)

Product Data



C57BL/6 mouse bone marrow cells were stained with CD11b (clone M1/70) APC and biotinylated CD64 (clone X54-5/7.1) (top) or mouse IgG1, κ isotype control (bottom) followed by Streptavidin-PE.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587