

PE anti-human CD8 Antibody

Catalog# / Size	344705 / 25 tests 344706 / 100 tests
Clone	SK1
Regulatory Status	RUO
Other Names	T8, Leu2
Isotype	Mouse IgG1, κ
Description	CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the $\alpha 3$ domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck.

Product Details

Verified Reactivity	Human, Cynomolgus, Rhesus
Reported Reactivity	African Green, Chimpanzee, Pigtailed Macaque, Sooty Mangabey
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Clone SK1 recognizes the a chain of CD8. Additional reported applications (for the relevant formats) include: proteogenomics ⁸ , immunohistochemistry of acetone-fixed frozen tissue sections, and spatial biology (IBEX) ^{9,10} . This clone was tested in-house and does not demonstrate utility for formalin-fixed paraffin-embedded (FFPE) human tonsil sections.
Application References	1. Ledbetter JA, <i>et al.</i> 1981. <i>J. Exp. Med.</i> 153:310. 2. Campanelli R, <i>et al.</i> 2002. <i>Intl. Immunol.</i> 14:39. 3. Evans RL, <i>et al.</i> 1981. <i>Immunol.</i> 78:544. 4. Wooldridge L, <i>et al.</i> 2005. <i>J. Bio. Chem.</i> 280:27491. 5. Ch'el IL, <i>et al.</i> 2011. <i>J Exp Med.</i> 208:633. PubMed 6. Carbone A, <i>et al.</i> 1999. <i>Blood</i> 93:2319. (IHC-F) 7. Ahmed A, <i>et al.</i> 2001. <i>J. Pathol.</i> 193:383. (IHC) 8. Peterson VM, <i>et al.</i> 2017. <i>Nat. Biotechnol.</i> 35:936. (PG) 9. Radtke AJ, <i>et al.</i> 2020. <i>Proc Natl Acad Sci USA.</i> 117:33455-33465. (SB) PubMed 10. Radtke AJ, <i>et al.</i> 2022. <i>Nat Protoc.</i> 17:378-401. (SB) PubMed
(PubMed link indicates BioLegend citation)	

Product Citations

1. Landry LG, *et al.* 2021. *Bio Protoc.* 11:e3883. [PubMed](#)
2. Zhang H, *et al.* 2021. *Cancer Gene Ther.* Online ahead of print. [PubMed](#)
3. Thaker YR, *et al.* 2022. *Front Oncol.* 12:884196. [PubMed](#)
4. Emmons TR, *et al.* 2021. *Cancer Immunol Res.* 9:790. [PubMed](#)
5. Crespo AC, *et al.* 2020. *Cell.* 182(5):1125-1139. [PubMed](#)
6. Vandooren J, *et al.* 2021. *Front Immunol.* 12:701739. [PubMed](#)
7. Massafra V, *et al.* 2021. *J Immunol.* 207:493. [PubMed](#)
8. Della-Torre E, *et al.* 2018. *Arthritis Rheumatol.* 70:1133. [PubMed](#)
9. Wang N, *et al.* 2021. *iScience.* 24:103205. [PubMed](#)
10. Price A, *et al.* 2016. *Proc Natl Acad Sci U S A.* 113: 10678 - 1068. [PubMed](#)
11. Leite NC, *et al.* 2020. *Cell Reports.* 32(2):107894.. [PubMed](#)
12. Meng W, *et al.* 2017. *Antimicrob Agents Chemother.* 10.1128/AAC.01719-17. [PubMed](#)
13. Meyer LK, *et al.* 2019. *J Clin Invest.* 130:863. [PubMed](#)
14. Ng CY, *et al.* 2022. *EMBO Rep.* 23:e54271. [PubMed](#)
15. Gimenez-Rivera V, *et al.* 2016. *J Immunol.* 197(11):4240-4246. [PubMed](#)
16. You L, *et al.* 2020. *Front Immunol.* 1.699305556. [PubMed](#)
17. Jing L, *et al.* 2020. *Cancer Immunol Res.* 8:648. [PubMed](#)
18. Zhong G, *et al.* 2022. *Oncoimmunology.* 11:2114740. [PubMed](#)
19. Pathania AS, *et al.* 2022. *Mol Ther Oncolytics.* 25:308. [PubMed](#)
20. Chen RH, *et al.* 2021. *Front Immunol.* 11:594775. [PubMed](#)
21. Amer HT, *et al.* 2022. *Front Mol Biosci.* 9:1032517. [PubMed](#)
22. Reuschl AK, *et al.* 2022. *Cell Rep.* 39:110650. [PubMed](#)
23. Liu Y, *et al.* 2018. *Cancer Cell.* 33:480. [PubMed](#)
24. Winkler C, *et al.* 2021. *JCI Insight.* 6:. [PubMed](#)
25. Ng CY, *et al.* 2022. *EMBO Rep.* 23:e55871. [PubMed](#)
26. Chen J, *et al.* 2021. *J Invest Dermatol.* 787:141. [PubMed](#)
27. Reitinger C, *et al.* 2022. *Front Immunol.* 13:970290. [PubMed](#)
28. Li Q, *et al.* 2022. *J Biol Chem.* :101753. [PubMed](#)
29. Wakeley ME, *et al.* 2020. *J Surg Res.* 245:610. [PubMed](#)
30. Lee IK, *et al.* 2020. *Cancers (Basel).* 12:. [PubMed](#)

RRID

AB_1953243 (BioLegend Cat. No. 344705)
AB_1953244 (BioLegend Cat. No. 344706)

Antigen Details

Structure	Ig superfamily, homodimer or heterodimer with CD8b, 32-34 kD
Distribution	Majority of thymocytes, T cell subset, NK cells
Function	MHC class I co-receptor, thymic differentiation, T cell activation
Ligand/Receptor	MHC Class I molecules
Cell Type	NK cells, T cells, Thymocytes
Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	1. Barclay N, <i>et al.</i> 1993. <i>The Leucocyte Antigen FactsBook.</i> Academic Press Inc. San Diego.
Gene ID	925

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Alexa Fluor® 647 anti-human CD8, Brilliant Violet 650™ anti-human CD8, Purified anti-human CD8, FITC anti-human CD8, PE anti-human CD8, PerCP anti-human CD8, PerCP/Cyanine5.5 anti-human CD8, PE/Cyanine7 anti-human CD8, APC/Cyanine7 anti-human CD8, Alexa Fluor® 488 anti-human CD8, Pacific Blue™ anti-human CD8, Biotin anti-human CD8, APC anti-human CD8, Alexa Fluor® 700 anti-human CD8, Purified anti-human CD8 (Maxpar® Ready), Brilliant Violet 510™ anti-human CD8, Brilliant Violet 711™ anti-human CD8, Brilliant Violet 785™ anti-human CD8, Brilliant Violet 605™ anti-human CD8, PE/Dazzle™ 594 anti-human CD8, APC/Fire™ 750 anti-human CD8, Brilliant Violet 421™ anti-human CD8, TotalSeq™-A0046 anti-human CD8, TotalSeq™-C0046 anti-human CD8, Brilliant Violet 750™ anti-human CD8, TotalSeq™-B0046 anti-human CD8, Spark Blue™ 550 anti-human

CD8, APC/Fire™ 810 anti-human CD8, PE/Fire™ 640 anti-human CD8, PE/Fire™ 700 anti-human CD8, TotalSeq™-D0046 anti-human CD8, GMP APC anti-human CD8, PE/Cyanine5 anti-human CD8 Antibody, Spark UV™ 387 anti-human CD8, GMP PE anti-human CD8, GMP PE/Cyanine7 anti-human CD8, Spark NIR™ 685 anti-human CD8, KIRAVIA Blue 520™ anti-human CD8, GMP FITC anti-human CD8, GMP Pacific Blue™ anti-human CD8, GMP PerCP anti-human CD8, Spark Violet™ 500 anti-human CD8

Product Data



Human peripheral blood lymphocytes stained with SK1 PE.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587