

## PE/Cyanine5 anti-human CD3 Antibody

|                          |   |
|--------------------------|---|
| <b>Catalog# / Size</b>   | 300410 / 100 tests  |
| <b>Clone</b>             | UCHT1   |
| <b>Regulatory Status</b> | RUO   |
| <b>Workshop</b>          | III 471   |
| <b>Other Names</b>       | T3, CD3ε  |
| <b>Isotype</b>           | Mouse IgG1, κ   |
| <b>Description</b>       | CD3ε is a 20 kD chain of the CD3/T-cell receptor (TCR) complex which is composed of two CD3ε, one CD3γ, one CD3δ, one CD3ζ (CD247), and a T-cell receptor (α/β or γ/δ) heterodimer. It is found on all mature T cells, NKT cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation. |

### Product Details

|   |   |
|---|---|
| <b>Verified Reactivity</b>                        | Human   |
| <b>Reported Reactivity</b>                        | Chimpanzee  |
| <b>Antibody Type</b>                              | Monoclonal  |
| <b>Host Species</b>                               | Mouse   |
| <b>Formulation</b>                                | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)   |
| <b>Preparation</b>                                | The antibody was purified by affinity chromatography, and conjugated with PE/Cyanine5 under optimal conditions.   |
| <b>Concentration</b>                              | Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)  |
| <b>Storage &amp; Handling</b>                     | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>   |
| <b>Application</b>                                | <a href="#">FC - Quality tested</a>   |
| <b>Recommended Usage</b>                          | Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.  |
| <b>Excitation Laser</b>                           | Blue Laser (488 nm)<br>Green Laser (532 nm)/Yellow-Green Laser (561 nm)   |
| <b>Application Notes</b>                          | Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections <sup>4,6,7</sup> and formalin-fixed paraffin-embedded sections <sup>11</sup> , immunoprecipitation <sup>1</sup> , activation of T cells <sup>2,3,5</sup> , Western blotting <sup>9</sup> , and spatial biology (IBEX) <sup>16,17</sup> . The LEAF™ purified antibody (Endotoxin < 0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 300413, 300414, and 300432). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 300437, 300438, 300465, 300466, 300473, 300474) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin < 0.01 EU/µg). |
| <b>Additional Product Notes</b>                   | BioLegend is in the process of converting the name PE/Cy5 to PE/Cyanine5. The dye molecule remains the same, so you should expect the same quality and performance from our PE/Cyanine5 products. Please contact <a href="#">Technical Service</a> if you have any questions.   |
| <b>Application References</b>                     | <ol style="list-style-type: none"> <li>Salmeron A, <i>et al.</i> 1991. <i>J. Immunol.</i> 147:3047. (IP)</li> <li>Graves J, <i>et al.</i> 1991. <i>J. Immunol.</i> 146:2102. (Activ)</li> <li>Lafont V, <i>et al.</i> 2000. <i>J. Biol. Chem.</i> 275:19282. (Activ)</li> <li>Ryschich E, <i>et al.</i> 2003. <i>Tissue Antigens</i> 62:48. (IHC)</li> </ol>  |
| <b>(PubMed link indicates BioLegend citation)</b> |   |

5. Thompson AG, *et al.* 2004. *J. Immunol.* 173:1671. (Activ)
6. Sakkas LI, *et al.* 1998. *Clin. Diagn. Lab. Immun.* 5:430. (IHC)
7. Mack CL, *et al.* 2004. *Pediatr. Res.* 56:79. (IHC)
8. Thakral D, *et al.* 2008. *J. Immunol.* 180:7431. (FC) [PubMed](#)
9. Van Dongen JMM, *et al.* 1988. *Blood* 71:603. (WB)
10. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
11. Pollard, K. *et al.* 1987. *J. Histochem. Cytochem.* 35:1329. (IHC)
12. Luckashenak N, *et al.* 2013. *J. Immunol.* 190:27. [PubMed](#)
13. Laurent AJ, *et al.* 2014. *PLoS One.* 9:103683. [PubMed](#)
14. Li J, *et al.* 2015. *Cancer Res.* 75:508. [PubMed](#)
15. Stoeckius M, *et al.* 2017. *Nat. Methods.* 14:865-868. (PG)
16. Radtke AJ, *et al.* 2020. *Proc Natl Acad Sci USA.* 117:33455-33465. (SB) [PubMed](#)
17. Radtke AJ, *et al.* 2022. *Nat Protoc.* 17:378-401. (SB) [PubMed](#)

#### Product Citations

1. Serra–Peinado C, *et al.* 2019. *Nat Commun.* 10:3705. [PubMed](#)
2. Evans RDR, *et al.* 2020. *Nat Commun.* 3.491666667. [PubMed](#)
3. Cassetta L *et al.* 2019. *Cancer Cell.* 35(4):588-602. [PubMed](#)
4. Jung J, *et al.* 2019. *Cell Rep.* 26:1906. [PubMed](#)
5. Orozco SL, *et al.* 2019. *Cell Rep.* 28:2275. [PubMed](#)

#### RRID

AB\_314064 (BioLegend Cat. No. 300410)

## Antigen Details

|                           |   |
|---------------------------|---|
| <b>Structure</b>          | Ig superfamily, with the subunits of CD3 $\gamma$ , CD3 $\delta$ , CD3 $\zeta$ (CD247) and TCR ( $\alpha/\beta$ or $\gamma/\delta$ ) forms CD3/TCR complex, 20 kD   |
| <b>Distribution</b>       | Mature T and NK T cells, thymocyte differentiation  |
| <b>Function</b>           | Antigen recognition, signal transduction, T cell activation   |
| <b>Ligand/Receptor</b>    | Peptide antigen bound to MHC  |
| <b>Cell Type</b>          | NKT cells, T cells, Thymocytes, Tregs   |
| <b>Biology Area</b>       | Immunology, Innate Immunity   |
| <b>Molecular Family</b>   | CD Molecules, TCRs  |
| <b>Antigen References</b> | <ol style="list-style-type: none"> <li>1. Barclay N, <i>et al.</i> 1993. <i>The Leucocyte FactsBook.</i> Academic Press. San Diego.</li> <li>2. Beverly P, <i>et al.</i> 1981. <i>Eur. J. Immunol.</i> 11:329.</li> <li>3. Lanier L, <i>et al.</i> 1986. <i>J. Immunol.</i> 137:2501-2507.</li> </ol> |
| <b>Gene ID</b>            | <a href="#">916</a>   |

## Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

APC anti-human CD3, Biotin anti-human CD3, FITC anti-human CD3, PE anti-human CD3, PE/Cyanine5 anti-human CD3, Purified anti-human CD3, Alexa Fluor® 647 anti-human CD3, Alexa Fluor® 488 anti-human CD3, Pacific Blue™ anti-human CD3, PE/Cyanine7 anti-human CD3, Alexa Fluor® 700 anti-human CD3, APC/Cyanine7 anti-human CD3, PerCP anti-human CD3, PerCP/Cyanine5.5 anti-human CD3, Brilliant Violet 421™ anti-human CD3, Brilliant Violet 570™ anti-human CD3, Ultra-LEAF™ Purified anti-human CD3, Purified anti-human CD3 (Maxpar® Ready), Alexa Fluor® 594 anti-human CD3, PE/Dazzle™ 594 anti-human CD3, Brilliant Violet 510™ anti-human CD3, Brilliant Violet 605™ anti-human CD3, Brilliant Violet 711™ anti-human CD3, Brilliant Violet 650™ anti-human CD3, APC/Fire™ 750 anti-human CD3, Brilliant Violet 785™ anti-human CD3, TotalSeq™-A0034 anti-human CD3, TotalSeq™-B0034 anti-human CD3, TotalSeq™-C0034 anti-human CD3, KIRAVIA Blue 520™ anti-human CD3, Spark Violet™ 538 anti-human CD3 Antibody, TotalSeq™-D0034 anti-human CD3, Spark Blue™ 574 anti-human CD3 Antibody, GMP Pacific Blue™ anti-human CD3, GMP PE anti-human CD3, GMP PE/Dazzle™ 594 anti-human CD3

## Product Data

Human peripheral blood lymphocytes  
stained with UCHT1 PE/Cyanine5



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587