

## TotalSeq™-A0857 anti-mouse CD34 Antibody

<b>Catalog# / Size</b>	128619 / 10 µg
<b>Clone</b>	HM34
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Mucosialin
<b>Isotype</b>	Armenian Hamster IgG
<b>Barcode Sequence</b>	GATTCCTTTACGAGC
<b>Description</b>	CD34 is a highly glycosylated hematopoietic progenitor antigen. Two isoforms of CD34 have been reported. CD34 is expressed on hematopoietic progenitors, as well as endothelial cells, brain and testis. CD34 is thought to function as an adhesion molecule for attachment of stem cells to extracellular matrix or stromal cells.

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Armenian Hamster
<b>Immunogen</b>	Mouse CD34 transfected BHK cells
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA.
<b>Preparation</b>	The antibody was purified by chromatography and conjugated with TotalSeq™-A oligomer under optimal conditions.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">PG - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> and the oligomer sequence is confirmed by sequencing. TotalSeq™-A antibodies are compatible with 10x Genomics Single Cell Gene Expression <a href="#">Solutions</a>.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
<b>Application Notes</b>	The HM34 antibody does not stain bone marrow cells like some other mouse CD34 antibodies, probably because the antibody recognizes a different epitope from other mAbs.
<b>Additional Product Notes</b>	<p>TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. <a href="#">10x Genomics Chromium System and Reagents</a>) and sequencer (e.g. Illumina analyzers) are required. Please contact <a href="#">technical support</a> for more information, or visit <a href="http://biolegend.com/totalseq">biolegend.com/totalseq</a>.</p> <p>The barcode flanking sequences are CCTTGGCACCCGAGAATTCCA (PCR handle), and BAAAAAAAAAAAAAAAAAAAAAAAAAAAAA*A*A (capture sequence). B represents either C, G, or T, and * indicates a phosphorothioated bond, to prevent nuclease degradation.</p>

View more applications data for this product in our [Scientific Poster Library](#).

## Product Citations

1. Guilliams M, *et al.* 2022. Cell. 185:379. [PubMed](#)

## RRID

AB\_2810392 (BioLegend Cat. No. 128619)

## Antigen Details

---

<b>Structure</b>	Type I membrane protein, highly glycosylated, 75-120 kd, two isoforms reported.
<b>Distribution</b>	Hematopoietic progenitors, brain, testis, endothelial cells
<b>Function</b>	Possible adhesion molecules thought to function in early hematopoiesis by mediating attachment of stem cells to bone marrow extracellular matrix or stromal cells. Present carbohydrate ligands to selectins.
<b>Ligand/Receptor</b>	L-selectin, and others.
<b>Cell Type</b>	Endothelial cells, Hematopoietic stem and progenitors
<b>Biology Area</b>	Cell Biology, Immunology, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	Adhesion Molecules, CD Molecules
<b>Antigen References</b>	1. Garlanda C, <i>et al.</i> 1997. <i>Eur J Cell Biol</i> 73:368 2. Brown J, <i>et al.</i> 1991. <i>Int Immunol</i> 3:175
<b>Gene ID</b>	<a href="#">12490</a>

## Related Protocols

---

[TotalSeq™-A Antibodies and Cell Hashing with 10x Single Cell 3' Reagent Kit v3 3.1 Protocol](#)

## Other Formats

---

Purified anti-mouse CD34, Biotin anti-mouse CD34, Alexa Fluor® 647 anti-mouse CD34, PerCP/Cyanine5.5 anti-mouse CD34, PE anti-mouse CD34, APC anti-mouse CD34, APC/Fire™ 750 anti-mouse CD34, PE/Dazzle™ 594 anti-mouse CD34, PE/Cyanine7 anti-mouse CD34, TotalSeq™-A0857 anti-mouse CD34, APC/Cyanine7 anti-mouse CD34, TotalSeq™-C0857 anti-mouse CD34, TotalSeq™-B0857 anti-mouse CD34

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587