

## TotalSeq™-C0109 anti-mouse CD16/32 Antibody

<b>Catalog# / Size</b>	101347 / 10 µg
<b>Clone</b>	93
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Fcγ R III/II, Ly-17
<b>Isotype</b>	Rat IgG2a, λ
<b>Barcode Sequence</b>	TTCGATGCTGGAGCA
<b>Description</b>	CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes and mediate adaptive immune responses.

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	Sorted pre-B cells
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
<b>Preparation</b>	The antibody was purified by chromatography and conjugated with TotalSeq™-C oligomer under optimal conditions.
<b>Concentration</b>	0.5 mg/mL
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
<b>Application</b>	<a href="#">PG - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> and the oligomer sequence is confirmed by sequencing. TotalSeq™-C antibodies are compatible with 10x Genomics Chromium Single Cell Immune Profiling <a href="#">Solution</a>.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
<b>Application Notes</b>	<p>Clone 93 can be used for blocking of CD16/CD32 interactions with the Fc domain of immunoglobulins, but is not the same clone as 2.4G2.</p> <p>The 93 mAb is specific to the common epitope of CD16/CD32. Additional reported applications (for the relevant formats) include: immunoprecipitation<sup>1</sup> and blocking of Fc-mediated reactions in functional studies<sup>2,4,23</sup>. It is useful for blocking non-specific binding of immunoglobulin to Fc receptors. For blocking of Fc receptors in flow cytometric analysis, pre-incubate the cells with purified anti-CD16/CD32 antibody (=1.0 µg per 10<sup>6</sup> cells in 100 µL volume) for 5-10 minutes on ice prior to immunostaining. For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 101330) (Endotoxin &lt;0.01 EU/µg, Azide-Free, 0.2 µm filtered).</p>

## Additional Product Notes

TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. [10x Genomics Chromium System and Reagents](#)) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit [biolegend.com/totalseq](http://biolegend.com/totalseq).

The barcode flanking sequences are CGGAGATGTGTATAAGAGACAGNNNNNNNNNN (PCR handle), and NNNNNNNNNCCCATATAAGA\*A\*A (capture sequence). N represents either randomly selected A, C, G, or T, and \* indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

## Application References

(PubMed link indicates BioLegend citation)

1. Personal communication (IP)
2. Oliver AM, *et al.* 1999. *Hybridoma* 18:113. (Block)
3. Brummel R and Lenert P. 2005. *J. Immunol.* 174:2429.
4. Terrazas LI, *et al.* 2005. *Int. J. Parasitol.* 35:1349. (Block)
5. Clements JL, *et al.* 2006. *J. Immunol.* 177:905.
6. Mohamed W, *et al.* 2010. *Infect Immun.* 78:3306. [PubMed](#)
7. Ouchi T, *et al.* 2011. *J. Exp Med.* 208:2607. [PubMed](#)
8. Kmiecik M, *et al.* 2011. *J. Vis. Exp.* 47:2381. [PubMed](#)
9. Yamazaki S, *et al.* 2012. *PLoS One.* 7:e51665. [PubMed](#)
10. Li J, *et al.* 2012. *Arthritis Rheum.* 64:1098. [PubMed](#)
11. Azuma M, *et al.* 2012. *Oncoimmunology.* 1:581. [PubMed](#)
12. Koon HW, *et al.* 2013. *J. Vis. Exp.* 68:4208. [PubMed](#)
13. Hegde VL, *et al.* 2013. *J Biol Chem.* 288:36810. [PubMed](#)
14. Huang J, *et al.* 2013. *J. Immunol Methods.* 387:254. [PubMed](#)
15. Dutow P, *et al.* 2014. *J Infect Dis.* [PubMed](#)
16. Fan Y, *et al.* 2014. *J Exp Med.* 211:313. [PubMed](#)
17. Huang HN, *et al.* 2014. *Antimicrob Agents Chemother.* 58:1538. [PubMed](#)
18. Takei S, *et al.* 2014. *Vaccine.* 32:3066. [PubMed](#)
19. Richardson ML, *et al.* 2014. *PLoS Negl Trop Dis.* 8:2825. [PubMed](#)
20. Cekanaviciute E, *et al.* 2014. *J Immunol.* 193:139. [PubMed](#)
21. Kimura T, *et al.* 2014. *Int Immunol.* 26:697. [PubMed](#)
22. Everad A, *et al.* 2014. *Nat Commun.* 5:5648. [PubMed](#)
23. Cenci E, *et al.* 2006. *J. Leuko. Biol.* 79(1):40-5. (Block)

## RRID

AB\_2819783 (BioLegend Cat. No. 101347)

## Antigen Details

---

<b>Structure</b>	Ig superfamily, 40-60 kD
<b>Distribution</b>	B cells, monocyte/macrophages, NK cells, neutrophils, mast cells, dendritic cells
<b>Function</b>	Low affinity receptors for IgG
<b>Ligand/Receptor</b>	IgG
<b>Cell Type</b>	B cells, Dendritic cells, Macrophages, Mast cells, Monocytes, Neutrophils, NK cells
<b>Biology Area</b>	Immunology, Innate Immunity
<b>Molecular Family</b>	CD Molecules, Fc Receptors
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Barclay AN, <i>et al.</i> 1997. <i>The Leukocyte Antigen FactsBook</i> Academic Press.</li><li>2. Unkeless JC. 1989. <i>J. Clin. Invest.</i> 83:355.</li><li>3. Qiu WQ, <i>et al.</i> 1990. <i>Science</i> 248:732.</li></ol>
<b>Gene ID</b>	<a href="#">14130</a> <a href="#">14131</a>

## Related Protocols

---

[TotalSeq™-B or -C with 10x Feature Barcoding Technology](#)

## Other Formats

---

Biotin anti-mouse CD16/32, FITC anti-mouse CD16/32, PE anti-mouse CD16/32, Purified anti-mouse CD16/32, Ultra-LEAF™ Purified anti-mouse CD16/32, Alexa Fluor® 647 anti-mouse CD16/32, PE/Cyanine7 anti-mouse CD16/32, TruStain FcX™ (anti-mouse CD16/32), PerCP/Cyanine5.5 anti-mouse CD16/32, APC anti-mouse CD16/32, APC/Cyanine7 anti-mouse CD16/32, Brilliant Violet 421™ anti-mouse CD16/32, Brilliant Violet 510™ anti-mouse CD16/32, Purified anti-mouse CD16/32 (Maxpar® Ready), Brilliant Violet 711™ anti-mouse CD16/32, TotalSeq™-A0109 anti-mouse CD16/32, TotalSeq™-B0109 anti-mouse CD16/32, TotalSeq™-C0109 anti-mouse CD16/32

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587