

## Ultra-LEAF™ Purified anti-human CD137 (4-1BB) Antibody

<b>Catalog# / Size</b>	309841 / 100 µg 309842 / 1 mg
<b>Clone</b>	4B4-1
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	VI C-7
<b>Other Names</b>	4-1BB, ILA, CD137, TNFRSF9
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	CD137 is a 39 kD transmembrane protein also known as 4-1BB. It is expressed on activated T cells. CD137 is a type I membrane protein and a member of the tumor necrosis factor receptor superfamily. CD137 appears to be important for T cell proliferation and survival, and induces monocyte activation through its interaction with 4-1BB ligand.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Reported Reactivity</b>	Chimpanzee, Baboon, Cynomolgus, Rhesus
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Ectodomain of recombinant human 4-1BB fusion protein
<b>Formulation</b>	0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.01 EU/µg of the protein (<0.001 ng/µg of the protein) as determined by the LAL test.
<b>Preparation</b>	The Ultra-LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.
<b>Concentration</b>	The antibody is bottled at the concentration indicated on the vial, typically between 2 mg/mL and 3 mg/mL. Older lots may have also been bottled at 1 mg/mL. To obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. This Ultra-LEAF™ solution contains no preservative; handle under aseptic conditions.
<b>Application</b>	<a href="#">FC - Quality tested</a> <a href="#">IP, FA, ELISA - Reported in the literature, not verified in house</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunoprecipitation <sup>1,4</sup> , inhibition of cytokine production <sup>2,3</sup> , and ELISA. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 309804) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated anti-mouse IgG second step (Cat. No. 405303), followed by Streptavidin-PE (Cat. No. 405204)).
<b>Application References</b>	1. Garni-Wagner B, <i>et al.</i> 1996. <i>Cell. Immunol.</i> 169:91. (IP) 2. Salih HR, <i>et al.</i> 2000. <i>J. Immunol.</i> 165:2903. (FA) 3. Kienzle G, <i>et al.</i> 2000. <i>Int. Immunol.</i> 12:73. (FA) 4. Langstein J, <i>et al.</i> 1998. <i>J. Immunol.</i> 160:2488. (IP)
<b>RRID</b>	AB_2810469 (BioLegend Cat. No. 309841) AB_2810470 (BioLegend Cat. No. 309842)

## Antigen Details

---

<b>Structure</b>	TNFR superfamily, type I transmembrane protein, 30 kD
<b>Distribution</b>	Activated T cells
<b>Function</b>	T cell costimulation
<b>Ligand/Receptor</b>	4-1BB ligand
<b>Cell Type</b>	T cells
<b>Biology Area</b>	Costimulatory Molecules, Immunology
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Gruss H, <i>et al.</i> 1995. <i>Blood</i> 85:3378.</li><li>2. Sica G, <i>et al.</i> 2000. <i>Adv. Exp. Med. Biol.</i> 465:355.</li><li>3. Alderson M, <i>et al.</i> 1994. <i>Eur. J. Immunol.</i> 24:2219.</li><li>4. Schwarz H, <i>et al.</i> 1996. <i>Blood</i> 87:2839.</li></ol>
<b>Gene ID</b>	<a href="#">3604</a>

## Related Protocols

---

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

---

Purified anti-human CD137 (4-1BB), PE anti-human CD137 (4-1BB), Biotin anti-human CD137 (4-1BB), PE/Cyanine5 anti-human CD137 (4-1BB), APC anti-human CD137 (4-1BB), PerCP/Cyanine5.5 anti-human CD137 (4-1BB), Alexa Fluor® 700 anti-human CD137 (4-1BB), PE/Cyanine7 anti-human CD137 (4-1BB), Brilliant Violet 421™ anti-human CD137 (4-1BB), APC/Cyanine7 anti-human CD137 (4-1BB), Brilliant Violet 605™ anti-human CD137 (4-1BB), Alexa Fluor® 647 anti-human CD137 (4-1BB), PE/Dazzle™ 594 anti-human CD137 (4-1BB), Brilliant Violet 650™ anti-human CD137 (4-1BB), Brilliant Violet 711™ anti-human CD137 (4-1BB), APC/Fire™ 750 anti-human CD137 (4-1BB), TotalSeq™-A0355 anti-human CD137 (4-1BB), TotalSeq™-B0355 anti-human CD137 (4-1BB), TotalSeq™-C0355 anti-human CD137 (4-1BB), Ultra-LEAF™ Purified anti-human CD137 (4-1BB), Brilliant Violet 750™ anti-human CD137 (4-1BB), TotalSeq™-D0355 anti-human CD137 (4-1BB)

## Product Data

---



PHA-stimulated (3 days) human peripheral blood mononuclear cells stained with LEAF™ purified 4B4-1, followed by biotinylated anti-mouse IgG and Sav-PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587