

Ultra-LEAF™ Purified anti-human CD274 (B7-H1, PD-L1) Antibody

Catalog# / Size	329715 / 100 µg 329716 / 1 mg 329745 / 5 mg 329746 / 25 mg 329747 / 50 mg 329748 / 100 mg
Clone	29E.2A3
Regulatory Status	RUO
Other Names	Programmed cell death ligand 1 (PD-L1), B7 homolog 1 (B7-H1)
Isotype	Mouse IgG2b, κ
Description	CD274, also known as PD-L1 and B7-H1, is type I transmembrane glycoprotein that serves as a ligand for CD279 (PD-1). This interaction is believed to regulate the balance between the stimulatory and inhibitory signals needed for responses to microbes and maintenance of self-tolerance. CD274 is involved in the costimulation of T cell proliferation and IL-10 and IFN-γ production in an IL-2-dependent and CD279-independent manner. Conflicting data has shown that CD274 can inhibit T cell proliferation and cytokine production, and alternatively, enhance T cell activation. Other studies suggest that CD274 may signal bidirectionally, raising interesting implications for its expression in a wide variety of cell types, including T and B cells, antigen-presenting cells, and nonhematopoietic cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Full length human PD-L1
Formulation	0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.01 EU/µg of the protein (<0.001 ng/µg of the protein) as determined by the LAL test.
Preparation	The Ultra-LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.
Concentration	The antibody is bottled at the concentration indicated on the vial, typically between 2 mg/mL and 3 mg/mL. Older lots may have also been bottled at 1 mg/mL. To obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. This Ultra-LEAF™ solution contains no preservative; handle under aseptic conditions.
Application	FC - Quality tested IHC, Block - Reported in the literature, not verified in house
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	Clone 29E.2A3 is reported to recognize an epitope on PD-L1 within the PD-L1-CD80 binding region ⁵ . Additional reported applications (for the relevant formats) include: blocking ¹⁻³ and immunohistochemical staining of acetone-fixed frozen sections ¹ . The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 329715, 329716, 329745 - 329748). It has been observed that clone 29E.2A3 is able to bind to Alexa Fluor® 700 antibody conjugates during multi-color immunofluorescent staining. This interaction can be resolved by sequentially

staining with the 29E.2A3 antibody first and then followed by the Alexa Fluor® 700 conjugate of interest.

Clone 29E.2A3 does not work in Western blot applications⁷.

Application References

1. Brown J, *et al.* 2003. *J. Immunol.* 170:1257. (FC, IHC, Block)
2. Radziejcz H, *et al.* 2007. *J. Virol.* 81:2545. (Block)
3. Nakamoto N, *et al.* 2009. *PLoS Pathog.* 5:e1000313. (Block)
4. Barsoum IB, *et al.* 2014. *Cancer Res.* 74:665. [PubMed](#)
5. Haile, S *et al.* 2013. *J. Immunol.* 191:2829.
6. RL M, *et al.* 2015. *PNAS.* 112:6506-6514. [PubMed](#)
7. Mahoney KM, *et al.* 2015. *Cancer Immunol. Res.* 3:1308.

Product Citations

1. Wang H, *et al.* 2020. *J Exp Clin Cancer Res.* 39:29. [PubMed](#)
2. Rieder SA, *et al.* 2020. *Cell Mol Immunol.* . [PubMed](#)
3. Baleeiro RB, *et al.* 2022. *Oncoimmunology.* 11:2080329. [PubMed](#)
4. Curnock AP, *et al.* 2021. *JCI Insight.* 6:. [PubMed](#)
5. Meng Q, *et al.* 2021. *J Immunother Cancer.* 9:. [PubMed](#)

RRID

AB_11149486 (BioLegend Cat. No. 329715)
AB_11149168 (BioLegend Cat. No. 329716)
AB_2783198 (BioLegend Cat. No. 329745)
AB_2783199 (BioLegend Cat. No. 329746)
AB_2783200 (BioLegend Cat. No. 329747)
AB_2783201 (BioLegend Cat. No. 329748)

Antigen Details

Distribution	T cells, B cells, NK cells, monocytes/macrophages, granulocytes and dendritic cells
Function	CD274 is involved in the costimulatory signal, essential for T lymphocyte proliferation and production of IL-10 and IFN- γ , in an IL-2-dependent and a PD-1-CD1-independent manner. Its interaction with PD-1-CD1 inhibits T-cell proliferation and cytokine production.
Ligand/Receptor	PD-1 (PDCD1)
Cell Type	B cells, Dendritic cells, Fibroblasts, Granulocytes, Macrophages, Monocytes, NK cells, T cells
Biology Area	Cancer Biomarkers, Costimulatory Molecules, Immunology
Molecular Family	Adhesion Molecules, CD Molecules, Immune Checkpoint Receptors
Antigen References	1. Sharpe A, <i>et al.</i> 2007. <i>Nat. Immunol.</i> 8:239.
Gene ID	29126


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD274 (B7-H1, PD-L1), Biotin anti-human CD274 (B7-H1, PD-L1), PE anti-human CD274 (B7-H1, PD-L1), APC anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 421™ anti-human CD274 (B7-H1, PD-L1), Ultra-LEAF™ Purified anti-human CD274 (B7-H1, PD-L1), PE/Cyanine7 anti-human CD274 (B7-H1, PD-L1), Purified anti-human CD274 (B7-H1, PD-L1) (Maxpar® Ready), Brilliant Violet 711™ anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 605™ anti-human CD274 (B7-H1, PD-L1), GolnVivo™ Purified anti-human CD274 (B7-H1, PD-L1), PE/Dazzle™ 594 anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 785™ anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 510™ anti-human CD274 (B7-H1, PD-L1), PerCP/Cyanine5.5 anti-human CD274 (B7-H1, PD-L1), Brilliant Violet 650™ anti-human CD274 (B7-H1, PD-L1), Alexa Fluor® 594 anti-human CD274 (B7-H1, PD-L1), TotalSeq™-A0007 anti-human CD274 (B7-H1, PD-L1), TotalSeq™-B0007 anti-human CD274 (B7-H1, PD-L1), TotalSeq™-C0007 anti-human CD274 (B7-H1, PD-L1), TotalSeq™-D0007 anti-human CD274 (B7-H1, PD-L1), PE/Fire™ 810 anti-human CD274 (B7-H1, PD-L1) Antibody, PE/Cyanine5 anti-human CD274 (B7-H1, PD-L1), Spark YG™ 570 anti-human CD274 (B7-H1, PD-L1)

Product Data



PHA-stimulated (3 days) human peripheral blood lymphocytes were stained with purified CD274 (clone 29E.2A3) (filled histogram) or purified mouse IgG2b, κ isotype control (open histogram), followed by anti-mouse IgG PE.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587