

APC anti-human CD177 Antibody

Catalog# / Size	315807 / 25 tests 315808 / 100 tests
Clone	MEM-166
Regulatory Status	RUO
Workshop	HCDM listed
Other Names	Neutrophil specific antigen 1, NB1, polycythemia rubra vera 1
Isotype	Mouse IgG1, κ
Description	CD177 is also known as neutrophil specific antigen 1, NB1, and polycythemia rubra vera 1. It is a member of the uPAR family and is a GPI-linked cell surface glycoprotein with a molecular weight of 60 kD. CD177 is expressed on granulocytes and bone marrow progenitors (early erythroblasts, megakaryocytes). It is thought to be involved in allogeneic and autoimmune responses to neutrophils.

Product Details

Verified Reactivity	Human, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Human granulocytes
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunoprecipitation, Western blotting ⁵ , and immunofluorescence ⁴ .
Application References	<ol style="list-style-type: none">1. Leucocyte Typing VII. Mason D, <i>et al.</i> Eds, 2002 Oxford University Press.2. von Vietinghoff S, <i>et al.</i> 2007. <i>Blood</i> 109:4487. PubMed3. Korkmaz B, <i>et al.</i> 2008. <i>J. Biol. Chem.</i> 283:35976. PubMed4. von Vietinghoff S, <i>et al.</i> 2007. <i>Blood</i> 109:4487. (IF)5. Jankowska AM, <i>et al.</i> 2011. <i>Haematologica.</i> 96:954. (WB)
(PubMed link indicates BioLegend citation)	
RRID	AB_2565213 (BioLegend Cat. No. 315807) AB_2565214 (BioLegend Cat. No. 315808)

Antigen Details

Structure	uPAR family, GPI-linked cell surface glycoprotein, 60 kD
Distribution	Granulocytes, bone marrow progenitors (early erythroblasts, megakaryocytes)
Function	Antigen involved in neutrophil allo- and autoimmunity, function unknown
Modification	Glycosylated
Cell Type	Granulocytes, Hematopoietic stem and progenitors, Neutrophils
Biology Area	Immunology
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Leukocyte Typing VII. Mason D, <i>et al.</i> (Eds.) Oxford University Press (2002) 2. Kissel K, <i>et al.</i> 2001. <i>Eur. J. Immunol.</i> 31:1301. 3. Lalezari P, <i>et al.</i> 1971. <i>J. Clin. Invest.</i> 50:1108. 4. Temerinac S, <i>et al.</i> 2000. <i>Blood</i> 95:2569.
Gene ID	57126


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD177, FITC anti-human CD177, PE anti-human CD177, APC anti-human CD177, APC/Cyanine7 anti-human CD177, TotalSeq™-A0382 anti-human CD177

Product Data



Human peripheral blood granulocytes were stained with CD177 (clone MEM-166) APC (filled histogram) or mouse IgG1, κ APC isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587