

APC/Cyanine7 anti-human CD83 Antibody

Catalog# / Size	305329 / 25 tests 305330 / 100 tests
Clone	HB15e
Regulatory Status	RUO
Other Names	HB15
Isotype	Mouse IgG1, κ
Description	CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member of the immunoglobulin superfamily, CD83 is expressed on a subset of dendritic cells, Langerhans cells, and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen presentation and/or lymphocyte activation, the precise function of this protein is unknown. CD83 is considered to be a useful marker for mature dendritic cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	African Green, Baboon, Pigtailed Macaque, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections ⁴ .
Additional Product Notes	BioLegend is in the process of converting the name APC/Cy7 to APC/Cyanine7. The dye molecule remains the same, so you should expect the same quality and performance from our APC/Cyanine7 products. Please contact Technical Service if you have any questions.
Application References (PubMed link indicates BioLegend citation)	<ol style="list-style-type: none">1. Knapp W, <i>et al.</i> 1989. Leucocyte Typing IV. Oxford University Press New York.2. Zhou L, <i>et al.</i> 1995. <i>J. Immunol.</i> 154:3821.3. Cao W, <i>et al.</i> 2005. <i>Biochem. J.</i> 385:85.4. Lore K, <i>et al.</i> 2002. <i>AIDS</i> 16:683. (IHC)5. Cho H, <i>et al.</i> 2007. <i>Physiol Genomics</i> doi:10.1152/physiolgenomics.00051.2006
Product Citations	<ol style="list-style-type: none">1. Walsh RE, <i>et al.</i> 2020. <i>MAbs.</i> 12:1764829. PubMed2. Das J, <i>et al.</i> 2021. <i>Med (N Y).</i> 2:1269. PubMed3. Jost M, <i>et al.</i> 2021. <i>eLife.</i> 10:00. PubMed
RRID	AB_2566392 (BioLegend Cat. No. 305329)

Antigen Details

Structure	Ig superfamily, single chain transmembrane glycoprotein, 43 kD
Distribution	Dendritic cells, Langerhan cells, activated B and T cells
Cell Type	B cells, Dendritic cells, Langerhans cells, T cells
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	<ol style="list-style-type: none"> 1. Kozlow E, <i>et al.</i> 1993. <i>Blood</i> 81:454. 2. Zhou L, <i>et al.</i> 1992. <i>J. Immunol.</i> 149:735. 3. Zhou L, <i>et al.</i> 1995. <i>Blood</i> 86:3295.
Gene ID	9308


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-human CD83, Biotin anti-human CD83, FITC anti-human CD83, PE anti-human CD83, PE/Cyanine5 anti-human CD83, Purified anti-human CD83, Alexa Fluor® 488 anti-human CD83, Alexa Fluor® 647 anti-human CD83, PerCP/Cyanine5.5 anti-human CD83, Brilliant Violet 421™ anti-human CD83, PE/Cyanine7 anti-human CD83, PE/Dazzle™ 594 anti-human CD83, APC/Cyanine7 anti-human CD83, Brilliant Violet 711™ anti-human CD83, APC/Fire™ 750 anti-human CD83, Brilliant Violet 605™ anti-human CD83, Brilliant Violet 785™ anti-human CD83, TotalSeq™-A0359 anti-human CD83, TotalSeq™-C0359 anti-human CD83, TotalSeq™-B0359 anti-human CD83, TotalSeq™-D0359 anti-human CD83

Product Data



Human monocyte-derived dendritic cells (induced with GM-CSF + IL-4 + TNF- α) were stained with CD83 (clone HB15e) APC/Cyanine7 (filled histogram) or mouse IgG1, κ APC/Cyanine7 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587