

APC/Fire™ 750 anti-human CD193 (CCR3) Antibody

Catalog# / Size	310726 / 100 tests
Clone	5E8
Regulatory Status	RUO
Other Names	CC CKR3, MIP1-alpha receptor like-2, eotaxin receptor, CD193, CCR3
Isotype	Mouse IgG2b, κ
Description	CD193, also known as CC-chemokine receptor 3 (CCR3), CC CKR3, MIP1-alpha receptor like-2, and eotaxin receptor, is a member of the G protein-coupled seven transmembrane receptors family. It binds to the CC chemokines eotaxin, eotaxin-2, and eotaxin-3 with high affinity. CCR3 has also been reported to bind RANTES, MCP-3, and MCP-4 with low affinity. CCR3 receptor is expressed on human eosinophils, basophils, mast cells, mononuclear phagocytes, platelets, CD34 ⁺ hematopoietic progenitor cells, Th2-like lymphocytes, and keratinocytes. CCR3 is thought to play a role in allergic diseases such as bronchial asthma and allergic rhinitis. CCR3 is a co-receptor for HIV-1 and HIV-2, and the binding of eotaxin with CCR3 has been shown to inhibit HIV infection in some cell types.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported applications (for the relevant formats) include: The 5E8 antibody is useful for immunofluorescent staining and flow cytometric analysis of CCR3 expression. It has been observed that the 5E8 antibody clone can interact with PE/Cy7 antibody conjugates during multi-color staining, potentially leading to unwanted staining. This interaction can be resolved by sequentially staining with the 5E8 antibody first and then followed by the PE/Cy7 conjugate of interest.
Application References	1. Beauvillian C, <i>et al.</i> 2011. <i>Blood</i> 117:1196. PubMed
(PubMed link indicates BioLegend citation)	
RRID	AB_2650970 (BioLegend Cat. No. 310726)


Antigen Details

Structure	G-protein coupled seven transmembrane domain receptor, 356 amino acids
Distribution	Eosinophils, basophils, mast, mononuclear phagocytes, platelets, CD34 ⁺ hematopoietic progenitor, Th2, keratinocytes
Function	Co-receptor for HIV-1 and HIV-2, allergy
Receptors	Eotaxin, eotaxin-2, eotaxin-3
Cell Type	Eosinophils, Erythrocytes, Hematopoietic stem and progenitors, Macrophages, Mast cells, Thymocytes
Biology Area	Immunology
Molecular Family	CD Molecules, Cytokine/Chemokine Receptors, GPCR
Antigen References	1. Gerard W, <i>et al.</i> 1996. <i>J. Exp. Med.</i> 183:2437. 2. Ugucioni C, <i>et al.</i> 1997. <i>J. Clin. Invest.</i> 100:1137. 3. Sallusto F, <i>et al.</i> 1997. <i>Science.</i> 277:2005. 4. Loetscher P, <i>et al.</i> 2001. <i>J. Biol. Chem.</i> 276:2986.
Regulation	Upregulated by high affinity Fc IgE receptor ligation (mast cells), RANTES (keratinocytes), IFNg (monocytes, neutrophils), HIV tat protein (basophils), IL-3, IL-5 and GM-CSF (CD34 ⁺ progenitor cells), IL-2 and IL-4(T cells). Downregulated by IL-
Gene ID	1232

Other Formats

Purified anti-human CD193 (CCR3), PE anti-human CD193 (CCR3), Brilliant Violet 605™ anti-human CD193 (CCR3), APC anti-human CD193 (CCR3), Alexa Fluor® 647 anti-human CD193 (CCR3), APC/Cyanine7 anti-human CD193 (CCR3), Brilliant Violet 421™ anti-human CD193 (CCR3), PerCP/Cyanine5.5 anti-human CD193 (CCR3), FITC anti-human CD193 (CCR3), Brilliant Violet 510™ anti-human CD193 (CCR3), Biotin anti-human CD193 (CCR3), APC/Fire™ 750 anti-human CD193 (CCR3), PE/Dazzle™ 594 anti-human CD193 (CCR3), TotalSeq™-A0397 anti-human CD193 (CCR3), TotalSeq™-C0397 anti-human CD193 (CCR3), Brilliant Violet 711™ anti-human CD193 (CCR3) Antibody, TotalSeq™-D0397 anti-human CD193 (CCR3)

Product Data



Human peripheral blood leukocytes were stained with CD16 FITC and CD193 (clone 5E8) APC/Fire™ 750 (top) or mouse IgG2b, κ APC/Fire™ 750 isotype control (bottom). Data shown is gated on granulocytes.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587