

Pacific Blue™ anti-mouse TCR β chain Antibody

Catalog# / Size	109225 / 25 µg 109226 / 100 µg
Clone	H57-597
Regulatory Status	RUO
Other Names	TCR-β chain, TCR-β, β-TCR
Isotype	Armenian Hamster IgG
Description	T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR-β is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR-α). It is expressed on α/β TCR-bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Armenian Hamster
Immunogen	Affinity purified TCR from mouse DO-11.10 cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Preparation	The antibody was purified by affinity chromatography, and conjugated with Pacific Blue™ under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . The suggested use of this reagent is ≤ 0.5 µg per 10 ⁶ cells in 100 µl volume or 100 µl of whole blood. It is highly recommended that the reagent be titrated for optimal performance for each application. * Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome. Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation. View full statement regarding label licenses
Excitation Laser	Violet Laser (405 nm)
Application Notes	H57-597 is a hamster mAb directed to an epitope of the C region of TCR β chain ¹² . The H57-597 antibody does not cross-react with γ/δ TCR-bearing T cells. Immobilized or soluble H57-597 antibody can activate α/β TCR-bearing T cells. Additional reported applications (for the relevant formats) for this antibody include: immunoprecipitation ² , <i>in vitro</i> stimulation ^{2,3} , <i>in vivo</i> depletion ⁴⁻⁶ , and immunohistochemical staining of acetone-fixed frozen sections ^{7,8,9} . The Ultra-LEAF™ purified antibody (Endotoxin <0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 109253-109258).
Application References	1. Gascoigne NJ. 1990. <i>J. Biol. Chem.</i> 265:9296. 2. Kruisbeek A, <i>et al.</i> 1991. <i>In Current Protocols in Immunology.</i> pp. 3.12.1. (Costim IP) 3. Davenport C, <i>et al.</i> 1995. <i>J. Immunol.</i> 155:3742. (Costim)
(PubMed link indicates BioLegend citation)	

4. Drobyski W, *et al.* 1996. *Blood* 87:5355. (Deplete)
5. Kummer U, *et al.* 2001. *Immunol. Lett.* 75:153. (Deplete)
6. van der Heyde HC, *et al.* 1995. *J. Immunol.* 154:3985. (Deplete)
7. Tomita K, *et al.* 1999. *Genes Dev.* 13:1203. (IHC)
8. Podd BS, *et al.* 2006. *J. Immunol.* 176:6532. (IHC)
9. Ponomarev ED, *et al.* 2007. *J. Immunol.* 178:39. (IHC)
10. Chappaz S, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-074245. (FC) [PubMed](#)
11. Tsukumo S, *et al.* 2006. *J. Immunol.* 177:8365. (FC) [PubMed](#)
12. Grégoire C, *et al.* 1991. *Proc. Natl. Acad. Sci USA* 88:8077.

Product Citations

1. Wu N, *et al.* 2020. *Cell Rep.* 30:1129. [PubMed](#)
2. Gilleron J *et al.* 2018. *Cell reports.* 25(12):3329-3341 . [PubMed](#)
3. Westerterp M *et al.* 2017. *Cell metabolism.* 25(6):1294-1304 . [PubMed](#)
4. Chen WS, *et al.* 2021. *Cell Rep.* 37:109974. [PubMed](#)
5. Schiller M, *et al.* 2021. *Immunity.* 54(5):1022-1036.e8. [PubMed](#)
6. Gan P, *et al.* 2017. *J Immunol.* 10.4049/jimmunol.1602025. [PubMed](#)
7. Dolina JS, *et al.* 2020. *Cell Rep.* 107249:31. [PubMed](#)
8. Lu Y, *et al.* 2021. *Gastroenterology.* 161:575. [PubMed](#)
9. Mintz MA, *et al.* 2019. *Immunity.* 51:310. [PubMed](#)
10. Sutherland TE, *et al.* 2021. *Frontiers in Immunology.* 12:715209. [PubMed](#)
11. La Rose AM, *et al.* 2021. *Mol Metab.* 53:101265. [PubMed](#)
12. Bankoti R, *et al.* 2017. *Sci Rep.* 10.1038/s41598-017-12171-3. [PubMed](#)
13. Pilzecker B, *et al.* 2016. *Nucleic Acids Res.* 44: 4734 - 4744. [PubMed](#)
14. Frost JN, *et al.* 2021. *Med (N Y).* 2:164. [PubMed](#)
15. Phalke SP, *et al.* 2019. *PLoS One.* 14:e0218827. [PubMed](#)
16. Dang EV *et al.* 2017. *Cell.* 171(5):1057-1071 . [PubMed](#)
17. Martina M, *et al.* 2016. *J Am Soc Nephrol.* 27: 1113-1123. [PubMed](#)
18. Redpath S, *et al.* 2015. *Infect Immun.* 83: 3881 - 3889. [PubMed](#)
19. Wang X, *et al.* 2022. *Elife.* 11:. [PubMed](#)
20. Wang Y, *et al.* 2022. *Nat Cell Biol.* 24:230. [PubMed](#)

RRID

AB_1027655 (BioLegend Cat. No. 109225)
 AB_1027649 (BioLegend Cat. No. 109226)

Antigen Details

Structure	Ig superfamily, CD3/TCR complex with CD3 and TCR α subunit
Distribution	Majority of T cells and thymocytes (correlated to differentiation)
Function	Antigen recognition, T cell activation
Ligand/Receptor	Peptide bound-MHC class I and II
Antigen References	<ol style="list-style-type: none"> 1. Davis MM, <i>et al.</i> 1998. <i>Ann. Rev. Immunol.</i> 16:523. 2. Huppa JB, <i>et al.</i> 2003. <i>Nat. Immunol.</i> 4:749. 3. Kubo R, <i>et al.</i> 1989. <i>J. Immunol.</i> 142:2736.
Gene ID	21577

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-mouse TCR β chain, Biotin anti-mouse TCR β chain, FITC anti-mouse TCR β chain, PE anti-mouse TCR β chain, PE/Cyanine5 anti-mouse TCR β chain, Purified anti-mouse TCR β chain, Alexa Fluor® 488 anti-mouse TCR β chain, Alexa Fluor® 647 anti-mouse TCR β chain, APC/Cyanine7 anti-mouse TCR β chain, PE/Cyanine7 anti-mouse TCR β chain, Alexa Fluor® 700 anti-mouse TCR β chain, Pacific Blue™ anti-mouse TCR β chain, Brilliant Violet 421™ anti-mouse TCR β chain, PerCP/Cyanine5.5 anti-mouse TCR β chain, Brilliant Violet 570™ anti-mouse TCR β chain, Brilliant Violet 510™ anti-mouse TCR β chain, Purified anti-mouse TCR β chain (Maxpar® Ready), Alexa Fluor® 594 anti-mouse TCR β chain, PE/Dazzle™ 594 anti-mouse TCR β chain, Brilliant Violet 605™ anti-mouse TCR β chain, Brilliant Violet 711™ anti-mouse TCR β chain, APC/Fire™ 750 anti-mouse TCR β chain, TotalSeq™-A0120 anti-mouse TCR β chain, Brilliant Violet 785™ anti-mouse TCR β chain, Brilliant Violet 650™ anti-mouse TCR β chain, Ultra-LEAF™ Purified anti-mouse TCR β chain, TotalSeq™-C0120 anti-mouse TCR β chain, TotalSeq™-B0120 anti-mouse TCR β chain

Product Data

C57BL/6 splenocytes stained with H57-597 Pacific Blue™



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587