

## Purified anti-human CD195 (CCR5) Antibody

<b>Catalog# / Size</b>	359101 / 25 µg 359102 / 100 µg
<b>Clone</b>	J418F1
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	C-C chemokine receptor type 5 (CCR5), HIV-1 fusion co-receptor
<b>Isotype</b>	Rat IgG2b, κ
<b>Description</b>	CD195, also known as CCR5, is a 45 kD G protein-coupled seven transmembrane CC-chemokine receptor. It binds to MIP-1α, MIP-1β, and RANTES and is expressed on a subset of T cells and monocytes. CCR5 mediates an intracellular signal thought to induce cell differentiation and proliferation. CCR5 has also been shown to act as a co-receptor for R5 HIV-1 cell entry; modification of CCR5 by sulfation contributes to the efficiency of HIV-1 entry. Studies have shown CCR5 to play a role in a variety of other human diseases, ranging from infectious and inflammatory diseases to cancer.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	Human CCR5 transfectants
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C.
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Product Citations</b>	<ol style="list-style-type: none"><li>1. Meckiff BJ, <i>et al.</i> 2019. J Immunol. 203:1276. <a href="#">PubMed</a></li><li>2. Yang R, <i>et al.</i> 2020. Cell. 183(7):1826-1847.e31. <a href="#">PubMed</a></li><li>3. Del Alcazar D, <i>et al.</i> 2019. Cell Rep. 28:3047. <a href="#">PubMed</a></li><li>4. Roussel M, <i>et al.</i> 2021. Cell Reports Medicine. 2(6):100291. <a href="#">PubMed</a></li></ol>
<b>RRID</b>	AB_2562456 (BioLegend Cat. No. 359101) AB_2562457 (BioLegend Cat. No. 359102)

### Antigen Details

---

<b>Structure</b>	G-coupled receptor family 1, membrane protein, 45 kD
<b>Distribution</b>	Subset of T cells, monocytes
<b>Function</b>	Binds C-C chemokines and transduces an intracellular signal thought to result in proliferation and differentiation, acts as a co-receptor with CD4 for HIV-1
<b>Ligand/Receptor</b>	MIP-1α, MIP-1β, RANTES

<b>Cell Type</b>	Monocytes, T cells
<b>Biology Area</b>	Cell Biology, Immunology, Innate Immunity, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	CD Molecules, Cytokine/Chemokine Receptors, GPCR
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Samson M, <i>et al.</i> 1996. <i>Biochemistry</i> 35:3362.</li> <li>2. Raport CJ, <i>et al.</i> 1996. <i>J. Biol. Chem.</i> 271:17161.</li> <li>3. Combadiere C, <i>et al.</i> 1996. <i>J. Leukoc. Biol.</i> 60:147.</li> <li>4. Deng H, <i>et al.</i> 1996. <i>Nature</i> 381:661.</li> <li>5. Lai J, <i>et al.</i> 2003. <i>CVI.</i> 10:1123.</li> <li>6. Mañes S, <i>et al.</i> 2003. <i>J. Exp. Med.</i> 198:1381.</li> <li>7. Vaday GG, <i>et al.</i> 2006. <i>Prostate</i> 66:124.</li> </ol>
<b>Gene ID</b>	<a href="#">1234</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-human CD195 (CCR5), Alexa Fluor® 488 anti-human CD195 (CCR5), PE anti-human CD195 (CCR5), PE/Cyanine7 anti-human CD195 (CCR5), APC/Cyanine7 anti-human CD195 (CCR5), PerCP/Cyanine5.5 anti-human CD195 (CCR5), Alexa Fluor® 647 anti-human CD195 (CCR5), Alexa Fluor® 700 anti-human CD195 (CCR5), Brilliant Violet 421™ anti-human CD195 (CCR5), FITC anti-human CD195 (CCR5), Pacific Blue™ anti-human CD195 (CCR5), PE/Dazzle™ 594 anti-human CD195 (CCR5), APC anti-human CD195 (CCR5), Brilliant Violet 510™ anti-human CD195 (CCR5), Brilliant Violet 711™ anti-human CD195 (CCR5), APC/Fire™ 750 anti-human CD195 (CCR5), Brilliant Violet 785™ anti-human CD195 (CCR5), TotalSeq™-A0141 anti-human CD195 (CCR5), TotalSeq™-C0141 anti-human CD195 (CCR5), TotalSeq™-B0141 anti-human CD195 (CCR5) Antibody, TotalSeq™-D0141 anti-human CD195 (CCR5), PE/Cyanine5 anti-human CD195 (CCR5)

## Product Data



Human peripheral blood lymphocytes were stained with purified CD195 (clone J418F1) (filled histogram), or rat IgG2b, κ (open histogram), followed by anti-rat IgG PE.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587