



Purified anti-mouse CD69 Recombinant Antibody

Catalog# / Size 164202 / 100 μg

Clone QA17A41

Regulatory Status RUO

Other Names Very Early Activation Antigen (VEA), AIM, EA1, MLR3, gp34/28

Isotype Rat lgG1, κ

Description CD69 is a 60 kD type II membrane protein composed of a 27/33 kD disulfide-linked homodimer,

also known as Very Early Activation (VEA) antigen, AlM, EA1, MLR3, and gp34/28. It is expressed on a subset of thymocytes and platelets. CD69 is rapidly induced on activated T and B cells, neutrophils, and NK cells. It is a C-type lectin, closely related to the NKR-P1 and Ly-49 NK cell activation molecules. CD69 is involved in the early events of cell activation and

thymocyte positive selection.

Product Details

Verified Reactivity Mouse

Antibody Type Recombinant

Host Species Rat

Formulation Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Preparation The antibody was purified by affinity chromatography.

Concentration 0.5 mg/mL

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C.

Application FC - Quality tested

Recommended Usage Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric

analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Product Citations
1. Ogbechi J, et al. 2022. Front Immunol. 13:1001956. PubMed

RRID AB_2904310 (BioLegend Cat. No. 164202)

Antigen Details

Structure C-type lectin, 27/33 kD

Distribution Activated T cells and B cells, NK cells, granulocytes, thymocytes, platelets

Function Lymphocyte activation

Cell Type B cells, Granulocytes, NK cells, Platelets, T cells, Thymocytes, Tregs

Biology Area Costimulatory Molecules, Immunology, Innate Immunity

Molecular Family CD Molecules

Gene ID <u>12515</u>


Related Protocols

Cell Surface Flow Cytometry Staining Protocol

Other Formats

Purified anti-mouse CD69 Recombinant Antibody, PE anti-mouse CD69 Recombinant Antibody

Product Data



PMA+ionomycin stimulated (6 hours) C57BL/6 mouse splenocytes stained with purified anti-mouse CD69 recombinant (clone QA17A41) (filled histogram) or purified rat lgG1, κ isotype control (open histogram), followed by goat anti-rat lgG PE.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering/#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587