

## TotalSeq™-C0135 anti-human CD324 (E-Cadherin) Antibody

<b>Catalog# / Size</b>	324127 / 10 µg
<b>Clone</b>	67A4
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	E-Cadherin, cadherin-1, and UVO
<b>Isotype</b>	Mouse IgG1, κ
<b>Barcode Sequence</b>	ATCCTTCTCCCTTC
<b>Description</b>	<p>The 67A4 antibody recognizes human CD324 also known as E-cadherin, cadherin-1, and UVO. CD324, a member of the cadherin superfamily, is a calcium-dependent, transmembrane cell-cell adhesion glycoprotein composed of 4 extracellular cadherin repeats and a highly conserved cytoplasmic tail region with a predicted molecular weight of approximately 100 kD. CD324 is widely expressed in epithelial cells in the colon, uterus, liver, keratinocytes, brain, heart, muscle, kidney, and pancreas, as well as erythroid cells. CD324 functions as a cell adhesion molecule involved in development, bacterial pathogenesis, and tumor invasion. In bacterial pathogenesis, the ectodomain of CD324 mediates bacterial adhesion to mammalian cells, while the cytoplasmic domain is required for internalization. CD324 binds to the αEβ7 integrin to mediate cell adhesion and also interacts with a number of intracellular proteins including including erbin, ezrin, caspase-3, caspase 8, β-catenin, presenilin 1, casein kinase II, as well as other extracellular proteins including the EGF receptor. CD324 is phosphorylated on multiple residues (S857, S866, S870, S872), and can be proteolytically cleaved at residue D769 by caspase-3. The 67A4 antibody has been shown to be useful for flow cytometry.</p>

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	T-47D cells
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
<b>Preparation</b>	The antibody was purified by chromatography and conjugated with TotalSeq™-C oligomer under optimal conditions.
<b>Concentration</b>	0.5 mg/mL
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">PG - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> and the oligomer sequence is confirmed by sequencing. TotalSeq™-C antibodies are compatible with 10x Genomics Chromium Single Cell Immune Profiling <a href="#">Solution</a> .

To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.

Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.

<b>Additional Product Notes</b>	TotalSeq™ reagents are designed to profile protein levels at a single cell level following an
---------------------------------	---

optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. [10x Genomics Chromium System and Reagents](#)) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit [biolegend.com/totalseq](#).

The barcode flanking sequences are CGGAGATGTGTATAAGAGACAGNNNNNNNNNN (PCR handle), and NNNNNNNNNCCCATATAAGA\*A\*A (capture sequence). N represents either randomly selected A, C, G, or T, and \* indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

## Application References

(PubMed link indicates BioLegend citation)

1. Armeanu S, *et al.* 1995. *J. Cell Biol.* 131:243.
2. Bühring HJ, *et al.* 1996. *Leukemia* 10:106.
3. Yauch RL, *et al.* 2005. *Clin. Cancer Res.* 11:8686. (WB)
4. Oetzuerk-Winder F, *et al.* 2012. *EMBO J.* 31:3431. (FC) [PubMed](#)
5. Ardehali R, *et al.* 2013. *PNAS.* 110:3405. [PubMed](#)
6. Rasanen K, *et al.* 2013. *Mol Cell Proteomics.* 12:3778. [PubMed](#)
7. Chaudhury A, *et al.* 2014. *Nucleic Acids Res.* 42:86. [PubMed](#)
8. Milne P, *et al.* 2015. *Blood.* 125:470. [PubMed](#)

**RRID** AB\_2832636 (BioLegend Cat. No. 324127)

## Antigen Details

<b>Structure</b>	Member of the cadherin superfamily. Calcium-dependent, transmembrane cell-cell adhesion glycoprotein composed of 4 extracellular cadherin repeats and a highly conserved cytoplasmic tail region. Predicted molecular weight approximately 100 kD.
<b>Distribution</b>	Widely expressed in epithelial cells in the colon, uterus, liver, keratinocytes, brain, heart, muscle, kidney, and pancreas, as well as erythroid cells
<b>Function</b>	Cell adhesion molecule involved in development, bacterial pathogenesis, and tumor invasion. The ectodomain of CD324 mediates bacterial adhesion to mammalian cell, while the cytoplasmic domain is required for internalization.
<b>Interaction</b>	Interacts with a variety of proteins including erbin, ezrin, caspase-3, caspase 8, EGF receptor, $\beta$ -catenin, presenilin 1, casein kinase II, and others
<b>Ligand/Receptor</b>	$\alpha\beta$ 7 integrin
<b>Modification</b>	Phosphorylated on multiple residues (S857, S866, S870, S872), proteolytically cleaved at residue D769 by caspase-3.
<b>Cell Type</b>	Embryonic Stem Cells, Epithelial cells, Erythrocytes
<b>Biology Area</b>	Cell Biology, Immunology, Neuroscience, Stem Cells, Synaptic Biology
<b>Molecular Family</b>	Adhesion Molecules, CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Overduin M, <i>et al.</i> 1995. <i>Science</i> 267:386.</li><li>2. Boggon TJ, <i>et al.</i> 2002. <i>Science</i> 296:1303.</li><li>3. Bex G, <i>et al.</i> 1995. <i>EMBO J.</i> 14:6107.</li><li>4. Perl AK, <i>et al.</i> 1998. <i>Nature</i> 392:190.</li></ol>
<b>Gene ID</b>	<a href="#">999</a>

## Related Protocols

[TotalSeq™-B or -C with 10x Feature Barcoding Technology](#)

## Other Formats

Purified anti-human CD324 (E-Cadherin), FITC anti-human CD324 (E-Cadherin), PE anti-human CD324 (E-Cadherin), APC anti-human CD324 (E-Cadherin), Alexa Fluor® 488 anti-human CD324 (E-Cadherin), Alexa Fluor® 647 anti-human CD324 (E-Cadherin), PerCP/Cyanine5.5 anti-human CD324 (E-Cadherin), PE/Cyanine7 anti-human CD324 (E-Cadherin), Alexa Fluor® 594 anti-human CD324 (E-Cadherin), APC/Fire™ 750 anti-human CD324 (E-Cadherin), Biotin anti-human CD324 (E-Cadherin), TotalSeq™-A0135 anti-human CD324 (E-Cadherin), TotalSeq™-C0135 anti-human CD324 (E-Cadherin), TotalSeq™-B0135 anti-human CD324 (E-Cadherin)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587