

## Alexa Fluor® 488 anti-mouse I-A<sup>b</sup> Antibody

<b>Catalog# / Size</b>	116410 / 100 µg
<b>Clone</b>	AF6-120.1
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	MHC class II
<b>Isotype</b>	Mouse (BALB/c) IgG2a, κ
<b>Description</b>	The AF6-120.1 antibody reacts with the I-A <sup>b</sup> MHC class II alloantigen. These class II molecules are expressed on antigen presenting cells (including B cells) and a subset of T cells from H-2 <sup>D</sup> bearing mice, and are involved in antigen presentation to T cells expressing CD3/TCR and CD4 proteins. The AF6-120.1 antibody cross-reacts with H-2 <sup>k</sup> and H-2 <sup>u</sup> haplotypes; this antibody does not cross-react with other haplotypes (d, f, q, r, s).

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	C57BL/10J splenocytes
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 488 under optimal conditions.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a>. For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.</p> <p>Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.</p> <p><a href="#">View full statement regarding label licenses</a></p>
<b>Application Notes</b>	<p>Additional reported applications (for relevant formats of this clone) include: immunohistochemical staining of frozen sections (acetone-fixed<sup>5</sup>; OCT-embedded, ethanol-fixed sections<sup>7</sup>), immunofluorescence microscopy<sup>3</sup> (including acetone-fixed epidermal sheets<sup>6</sup>), immunoprecipitation<sup>7,8</sup>. Directly conjugated antibody was used for IF in (3) and (6) and for IHC in (5).</p> <p>Does not react with other haplotypes (e.g., d, f, q, r, s).</p>
<b>Application References</b>	<ol style="list-style-type: none"> <li>1. Wall KA, <i>et al.</i> 1983. <i>J. Immunol.</i> 131:1056. (FC)</li> <li>2. Cohn LE, <i>et al.</i> 1986. <i>P. Natl. Acad. Sci. USA</i> 83:747. (FC)</li> <li>3. Inaba K, <i>et al.</i> 1998. <i>J. Exp. Med.</i> 188:2163 (IF)</li> <li>4. Hamrah P, <i>et al.</i> 2002. <i>Invest Ophthalmol Vis. Sci.</i> 43:639 (IF)</li> <li>5. Buono C, <i>et al.</i> 2003. <i>Arterioscler. Thromb. Vasc. Biol.</i> 23:454. (IHC)</li> <li>6. Wang Z, <i>et al.</i> 2004. <i>J. Immunol.</i> 172:5924. (IHC IF)</li> <li>7. Nakagawa TY, <i>et al.</i> 1999. <i>Immunity</i> 10:207. (IP)</li> <li>8. Podolin PL, <i>et al.</i> 2008. <i>J. Immunol.</i> 180:7989. (FC IP) <a href="#">PubMed</a></li> </ol>
<b>(PubMed link indicates BioLegend citation)</b>	

9. Schneppenheim J, et al. 2013. *J Exp Med.* 210:41. [PubMed](#).

## Product Citations

1. Shukla A, et al. 2021. *Adv Nanobiomed Res.* 1: [PubMed](#)
2. Caronni N, et al. 2018. *Cancer Res.* 78:1685. [PubMed](#)
3. Khare P, et al. 2017. *J Autoimmun.* 10.1016/j.jaut.2017.09.002. [PubMed](#)
4. Das A, et al. 2020. *J Bone Miner Res.* 36:199. [PubMed](#)

## RRID

AB\_493140 (BioLegend Cat. No. 116410)

## Antigen Details

---

<b>Structure</b>	MHC class II
<b>Distribution</b>	B cell and activated T cells, APCs of H-2 <sup>b</sup> mice
<b>Function</b>	Antigen presentation
<b>Ligand/Receptor</b>	CD3/TCR, CD4
<b>Cell Type</b>	Antigen-presenting cells, B cells, T cells
<b>Biology Area</b>	Immunology, Innate Immunity
<b>Molecular Family</b>	MHC Antigens
<b>Antigen References</b>	1. Watts C. 1997. <i>Annu. Rev. Immunol.</i> 15:821. 2. Pamer E, et al. 1998. <i>Annu. Rev. Immunol.</i> 16:323.
<b>Gene ID</b>	<a href="#">14961</a>

## Related Protocols

---

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

---

Biotin anti-mouse I-A<sup>b</sup>, FITC anti-mouse I-A<sup>b</sup>, PE anti-mouse I-A<sup>b</sup>, Purified anti-mouse I-A<sup>b</sup>, Alexa Fluor® 488 anti-mouse I-A<sup>b</sup>, Alexa Fluor® 647 anti-mouse I-A<sup>b</sup>, PerCP/Cyanine5.5 anti-mouse I-A<sup>b</sup>, APC anti-mouse I-A<sup>b</sup>, PE/Cyanine7 anti-mouse I-A<sup>b</sup>, Pacific Blue™ anti-mouse I-A<sup>b</sup>, APC/Fire™ 750 anti-mouse I-A<sup>b</sup>, APC/Cyanine7 anti-mouse I-A<sup>b</sup>

## Product Data

---



C57BL/6 mouse splenocytes stained with AF6-120.1 Alexa Fluor® 488

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic,

therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587