

## FITC anti-mouse F4/80 Recombinant Antibody

<b>Catalog# / Size</b>	157309 / 25 µg 157310 / 100 µg
<b>Clone</b>	QA17A29
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	EMR1, Ly71
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	F4/80 is a 160 kD glycoprotein. It is characterized as a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on the majority of tissue macrophages including peritoneal macrophages, macrophages in lung, gut, thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen, lymph node and Peyer's patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been identified, but it has been reported that F4/80 is required for induction of CD8 <sup>+</sup> T cells-mediated peripheral tolerance.

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Recombinant
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Murine macrophages
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions.
<b>Concentration</b>	0.5 mg/mL
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per million cells in 100 µL volume. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Excitation Laser</b>	Blue Laser (488 nm)
<b>Product Citations</b>	1. Kakizaki M, <i>et al.</i> 2021. Cell Death Dis. 12:1010. <a href="#">PubMed</a>
<b>RRID</b>	AB_2876535 (BioLegend Cat. No. 157309) AB_2876535 (BioLegend Cat. No. 157310)

### Antigen Details

---

<b>Structure</b>	A 160 kD glycoprotein and a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family
<b>Distribution</b>	Expressed on the majority of tissue macrophages

**Function** Required for induction of CD8<sup>+</sup> T cells-mediated peripheral tolerance

**Cell Type** Macrophages, Monocytes

**Biology Area** Immunology, Innate Immunity

#### Antigen References

1. Schaller E, *et al.* 2002. *Mol. Cell. Biol.* 22:8035.
2. Stevceva L, *et al.* 2001. *BMC Clin Pathol.* 1:3.
3. Kobayashi M, *et al.* 2008. *J. Leukoc. Biol.* 83:1354.
4. Poeckel D, *et al.* 2009. *J. Biol Chem.* 284:21077.
5. Glass AM, *et al.* 2013. *J. Immunol.* 190:4830.
6. Koehm S, *et al.* 2007. *J. Allergy Clin. Immunol.* 120:570.
7. Rankin AL, *et al.* 2010. *J. Immunol.* 184:1526.
8. Sasi SP, *et al.* 2014. *J Biol Chem.* 289:14178.
9. Thakus VS, *et al.* 2014. *Toxicol Lett.* 230:322.
10. Ohnishi A, *et al.* 2015. *PLoS One.* 10:117616.

**Gene ID** [13733](#)


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-mouse F4/80 Recombinant Antibody, PE anti-mouse F4/80 Recombinant Antibody, APC anti-mouse F4/80 Recombinant Antibody, PE/Cyanine7 anti-mouse F4/80 Recombinant Antibody, FITC anti-mouse F4/80 Recombinant Antibody, Spark YG™ 593 anti-mouse F4/80 Recombinant Antibody, APC/Cyanine7 anti-mouse F4/80 Recombinant Antibody, PerCP/Cyanine5.5 anti-mouse F4/80 Recombinant Antibody, Alexa Fluor® 647 anti-mouse F4/80 Recombinant Antibody, PE/Fire™ 640 anti-mouse F4/80 Recombinant Antibody

## Product Data



Thioglycollate-elicited Balb/c mouse peritoneal macrophages were stained with anti-mouse CD11b APC and anti-mouse F4/80 FITC (clone QA17A29) (left) or anti-mouse IgG1, κ isotype control FITC (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587