

PE anti-human CD80 Antibody

Catalog# / Size	305207 / 25 tests 305208 / 100 tests
Clone	2D10
Regulatory Status	RUO
Workshop	VI CD80.1
Other Names	B7-1, B7, BB1
Isotype	Mouse IgG1, κ
Description	CD80, also known as B7-1, B7, and BB1, is a 60 kD single chain type I glycoprotein belonging to the immunoglobulin superfamily. CD80 is expressed on activated B and T cells, macrophages, and dendritic cells. CD80 binds to CD28 and CD152 (CTLA-4). Along with CD86, CD80 plays a critical role in regulation of T cell activation. The interaction of CD80 with CD28 provides a potent costimulatory signal for T cell activation through the CD3 complex, while its interaction with CTLA-4 provides an inhibitory signal for T cell activation.

Product Details

Verified Reactivity	Human
Reported Reactivity	Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Additional reported applications (for the relevant formats) include: <i>in vitro</i> blocking of T cell activation, immunohistochemical staining of acetone-fixed frozen tissue sections ² , immunoprecipitation, and Western blotting ³ . The Ultra-LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. Nos. 305245 & 305246).
Application References	1. Kishimoto T, <i>et al.</i> Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London. 2. Battifora M. 1998. <i>J. Clin. Endocr. Metab.</i> 83:4130. (IHC) 3. Van der Merwe PA, <i>et al.</i> 1997. <i>J. Exp. Med.</i> 185:3. (WB) 4. Jayakumar A, <i>et al.</i> 2008. <i>Infect. Immun.</i> 76:2138. PubMed 5. Schubert DA, <i>et al.</i> 2012. <i>J. Exp. Med.</i> 209:335. PubMed 6. Wen T, <i>et al.</i> 2014. <i>J. Immunol.</i> 192:5481. PubMed
(PubMed link indicates BioLegend citation)	
Product Citations	1. Werz O, <i>et al.</i> 2018. <i>Nat Commun.</i> 9:59. PubMed

2. Lathrop S, *et al.* 2015. *Infect Immun* . 83: 2661 - 2671. [PubMed](#)
3. Li H, *et al.* 2016. *J Immunol*. 196: 4064 - 4074. [PubMed](#)
4. Hagan T, *et al.* 2020. *Cell*. 178(6):1313-1328.e13.. [PubMed](#)
5. Sugawara E, *et al.* 2020. *Autophagy*. 1.323611111. [PubMed](#)
6. Roskopf S, *et al.* 2016. *Sci Rep*. 6:31580. [PubMed](#)
7. Hou ZH, *et al.* 2020. *Am J Physiol Cell Physiol*. 318:C649. [PubMed](#)
8. Santiago K, *et al.* 2016. *Pathog Dis*. 74: ftw081. [PubMed](#)
9. Du X, *et al.* 2018. *Cell Res*. 28:416. [PubMed](#)
10. Guo Z, *et al.* 2022. *NPJ Precis Oncol*. 6:34. [PubMed](#)
11. Herath S, *et al.* 2014. *PLoS One*. 9:88327. [PubMed](#)
12. Wang D, *et al.* 2016. *J Cell Sci*. 129: 2261 - 2272. [PubMed](#)
13. Zhang S, *et al.* 2020. *Front Immunol*. 1.757638889. [PubMed](#)
14. Jutz S, *et al.* 2016. *J Immunol Methods*. 430:10-20. [PubMed](#)
15. Pellegrini JM, *et al.* 2021. *Sci Rep*. 11:13559. [PubMed](#)
16. Lewis SA, *et al.* 2021. *Front Immunol*. 12:724015. [PubMed](#)
17. Balan I, *et al.* 2022. *Front Immunol*. 13:940095. [PubMed](#)
18. Ganesan R, *et al.* 2020. *FASEB J*. 34:15888. [PubMed](#)
19. Zhang L, *et al.* 2011. *Blood*. 117:6184. [PubMed](#)
20. Blaszcak AM, *et al.* 2019. *J Diabetes Res*. 2019:8124563. [PubMed](#)
21. Serhan J 2012. *Blood*. 120:e60. [PubMed](#)
22. Nowak W, *et al.* 2020. *EBioMedicine*. 50:290-305.. [PubMed](#)
23. Luque-Martin R, *et al.* 2021. *J Immunol*. 207:555. [PubMed](#)
24. Zhao Y, *et al.* 2020. *Immunity*. 51(6):1059-1073.e9.. [PubMed](#)
25. Pi J, *et al.* 2022. *J Nanobiotechnology*. 20:36. [PubMed](#)
26. Kubala MH, *et al.* 2018. *Cell Rep*. 25:2177. [PubMed](#)
27. Gärtner K, *et al.* 2019. *J Extracell Vesicles*. 8:1573051. [PubMed](#)
28. Longbrake EE, *et al.* 2018. *Mult Scler*. 24:728. [PubMed](#)
29. Corgnac S, *et al.* 2022. *J Immunother Cancer*. 10:. [PubMed](#)
30. Chen R, *et al.* 2021. *Front Oncol*. 11:743050. [PubMed](#)
31. Morell M, *et al.* 2022. *EBioMedicine*. 76:103808. [PubMed](#)
32. Trabanelli S, *et al.* 2014. *J Immunol*. 192:1231. [PubMed](#)

RRID AB_314503 (BioLegend Cat. No. 305207)
 AB_314504 (BioLegend Cat. No. 305208)

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, 60 kD
Distribution	Activated B cells and T cells, macrophages, and dendritic cells
Function	T cell costimulation
Ligand/Receptor	CD28, CD152 (CTLA-4)
Cell Type	B cells, Dendritic cells, Macrophages, T cells, Tregs
Biology Area	Cell Biology, Costimulatory Molecules, Immunology, Neuroscience, Neuroscience Cell Markers
Molecular Family	CD Molecules, Immune Checkpoint Receptors
Antigen References	<ol style="list-style-type: none"> 1. Freeman G, <i>et al.</i> 1991. <i>J. Exp. Med.</i> 174:625. 2. Linsley P, <i>et al.</i> 1996. <i>Immunity</i> 4:535. 3. Linsley P, <i>et al.</i> 1991. <i>J. Exp. Med.</i> 174:561.
Gene ID	941

Related Protocols


[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Biotin anti-human CD80, FITC anti-human CD80, PE anti-human CD80, PE/Cyanine5 anti-human CD80, Purified anti-human CD80, Alexa Fluor® 488 anti-human CD80, Alexa Fluor® 647 anti-human CD80, PE/Cyanine7 anti-human CD80, APC anti-human CD80, Brilliant Violet 650™ anti-human CD80, Brilliant Violet 421™ anti-human CD80, Brilliant Violet 605™ anti-human CD80, PE/Dazzle™ 594 anti-human CD80, PerCP/Cyanine5.5 anti-human CD80, Brilliant Violet 510™ anti-human CD80, Brilliant Violet 711™ anti-human CD80, Brilliant Violet 785™ anti-human CD80, TotalSeq™-A0005 anti-human CD80, TotalSeq™-B0005 anti-human CD80,

TotalSeq™-C0005 anti-human CD80, Ultra-LEAF™ Purified anti-human CD80, Brilliant Violet 750™ anti-human CD80, TotalSeq™-D0005 anti-human CD80

Product Data



Human B-cell Burkitt's lymphoma cell line
Daudi stained with 2D10 PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587