

PE/Cyanine7 anti-IRF4 Antibody

| | |
|--------------------------|---|
| Catalog# / Size | 646413 / 25 µg 646414 / 100 µg |
| Clone | IRF4.3E4 |
| Regulatory Status | RUO |
| Other Names | Interferon regulatory factor 4 |
| Isotype | Rat IgG1, κ |
| Description | The IRF family consists of at least nine members. IRF4 and IRF8 are highly homologous to each other and also redundant in function. IRF4 is critical for Th2 and Th17 development. Together with TRF8, it plays an essential role in macrophage and dendritic cell development and function. IRF4 is also reported to be essential for pre-B cell development, receptor editing, germinal center reactor and plasma cell differentiation. |

Product Details

| | |
|---------------------------------|---|
| Verified Reactivity | Mouse, Human |
| Antibody Type | Monoclonal |
| Host Species | Rat |
| Immunogen | GST fusion protein containing C-terminal of murine IRF4 |
| Formulation | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide. |
| Preparation | The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under optimal conditions. |
| Concentration | 0.2 mg/ml |
| Storage & Handling | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze. |
| Application | ICFC - Quality tested |
| Recommended Usage | Each lot of this antibody is quality control tested by intracellular flow cytometry using our True-Nuclear™ Transcription Factor Staining Protocol . For flow cytometric staining, the suggested use of this reagent is =0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application. |
| Excitation Laser | Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm) |
| Application Notes | Additional reported applications (for the relevant formats) include: spatial biology (IBEX) ^{3,4} . |
| Additional Product Notes | BioLegend is in the process of converting the name PE/Cy7 to PE/Cyanine7. The dye molecule remains the same, so you should expect the same quality and performance from our PE/Cyanine7 products. Please contact Technical Service if you have any questions. |
| Application References | <ol style="list-style-type: none">1. Zheng Y, <i>et al.</i> 2009. <i>Nature</i> 458:3512. Yin SY, <i>et al.</i> 2011. <i>Exp Cell Res.</i> 317:2210. PubMed3. Radtke AJ, <i>et al.</i> 2020. <i>Proc Natl Acad Sci USA.</i> 117:33455-33465. (SB) PubMed4. Radtke AJ, <i>et al.</i> 2022. <i>Nat Protoc.</i> 17:378-401. (SB) PubMed |
| RRID | AB_2728479 (BioLegend Cat. No. 646413) AB_2728480 (BioLegend Cat. No. 646414) |

Antigen Details

| | |
|---------------------------|--|
| Structure | Molecular weight is approximately 51 kD. |
| Distribution | IRF4 is reported to be expressed exclusively in the immune system. |
| Biology Area | Cell Biology, Transcription Factors |
| Molecular Family | Nuclear Markers |
| Antigen References | 1. Lu R. 2008. <i>Trends Immunol</i> 29:487. |
| Gene ID | 3662 |

Related Protocols


[True-Nuclear™ Transcription Factor Staining Protocol for 96-Well U Bottom Plate](#)

[True-Nuclear™ Transcription Factor Staining Protocol for 5mL Tubes](#)

Other Formats

Alexa Fluor® 488 anti-IRF4, Purified anti-IRF4, PE anti-IRF4, Alexa Fluor® 647 anti-IRF4, Go-ChIP-Grade™ Purified anti-IRF4, PE/Cyanine7 anti-IRF4, PerCP/Cyanine5.5 anti-IRF4, Alexa Fluor® 594 anti-IRF4, Pacific Blue™ anti-IRF4

Product Data



PHA-stimulated (3 days) or freshly isolated human peripheral blood lymphocytes were stained with CD4 APC, this fixed and permeabilized with True-Nuclear™ Transcription Factor Buffer Set (Cat. No. 424401). Cells were then stained with IRF4 (clone IRF4.3E4) PE/Cyanine7 (open histogram with solid line for fresh cells, filled histogram for stimulated cells) or rat IgG1, ? PE/Cyanine7 isotype control (dash line for fresh cell, dot line for stimulated cells). Data shown was gated on CD4⁺ lymphocyte population.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587