

TotalSeq™-A0366 anti-human CD184 (CXCR4) Antibody

Catalog# / Size	306531 / 10 µg
Clone	12G5
Regulatory Status	RUO
Workshop	VII 70204
Other Names	CXCR4, Fusin
Isotype	Mouse IgG2a, κ
Barcode Sequence	TCAGGTCCTTTCAAC
Description	CD184, also known as fusin or CXCR4, is a 45 kD seven transmembrane G-protein-linked CXC chemokine receptor. CD184 is widely expressed on blood and tissue cells, including B and T cells, monocytes, macrophages, dendritic cells, granulocytes, megakaryocytes/platelets, lymphoid, myeloid precursor cells, endothelial cells, epithelial cells, astrocytes, and neurons, among other tissue cells. CD184 is the receptor for CXC chemokine SDF-1, mediates blood cell migration, and is involved in B lymphopoiesis and myelopoiesis, cardiogenesis, blood vessel formation, and cerebellar development. CXCR4 is also a coreceptor of X4 HIV-1 and an alternative receptor for some isolates of HIV-2.

Product Details

Verified Reactivity	Human, Cynomolgus, Rhesus
Reported Reactivity	African Green, Baboon, Chimpanzee, Sooty Mangabey
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	CP-MAC-infected Sup-T1 cells
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA.
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-A oligomer under optimal conditions.
Concentration	0.5 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-A antibodies are compatible with 10x Genomics Single Cell Gene Expression Solutions.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Application Notes	Additional reported applications (for the relevant formats) include: immunohistochemical staining of paraffin-embedded tissue sections ¹¹ , immunocytochemistry ³ , immunofluorescence microscopy ^{2,6} , and blocking of CD4-independent infection by HIV-2 and CD4-dependent infection by some T cell-

tropic isolates of HIV-14,5. Clone 12G5 may not be suitable for Western blotting.¹⁰ The Ultra-LEAF™ purified antibody (Endotoxin <0.01 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. Nos. 306539 & 306540).

Additional Product Notes

TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. [10x Genomics Chromium System and Reagents](#)) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit [biolegend.com/totalseq](#).

The barcode flanking sequences are CCTTGGCACCCGAGAATTCCA (PCR handle), and BAAA*A*A (capture sequence). B represents either C, G, or T, and * indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

Application References

(PubMed link indicates BioLegend citation)

1. McKnight A, *et al.* 1997. *J. Virol.* 71:1692.
2. Endres MJ, *et al.* 1996. *Cell* 87:745. (Immunogen, IF)
3. Volin MV, *et al.* 1998. *Biochem. Biophys. Res. Commun.* 242:46. (ICC)
4. Berndt C, *et al.* 1998. *P. Natl. Acad. Sci. USA* 95:12556. (Block)
5. Ullrich CK, *et al.* 2000. *Blood* 96:1438. (Block)
6. Murga M, *et al.* 2005. *Blood* 105:1992. (IF)
7. Thompson BD. 2007. *J. Biol. Chem.* 282:9547. (FC) [PubMed](#)
8. Isnardi I, *et al.* 2010. *Blood* 115:5026. [PubMed](#)
9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
10. Fischer T, *et al.* 2008. *PLoS One* 3:e4069.
11. Schmid BC, *et al.* 2004. *Breast Cancer Res. Treat.* 84:247. (IHC)

Product Citations

1. Hao Y, *et al.* 2021. *Cell.* 184:3573. [PubMed](#)

RRID

AB_2800790 (BioLegend Cat. No. 306531)

Antigen Details

Structure	Rhodopsin family, G-protein linked seven transmembrane glycoprotein, 45 kD
Distribution	T cells and B cells, dendritic cells, monocytes, granulocytes, hematopoietic progenitors, endothelial cells
Function	B lymphopoiesis and myelopoiesis, cardiogenesis, blood vessel formation, cerebellar development
Ligand/Receptor	SDF-1 receptor, coreceptor for X4 HIV-1
Cell Type	B cells, Dendritic cells, Endothelial cells, Granulocytes, Hematopoietic stem and progenitors, Mesenchymal Stem Cells, Monocytes, Neural Stem Cells, T cells, Tregs
Biology Area	Cell Biology, Immunology, Innate Immunity, Neuroinflammation, Neuroscience, Neuroscience Cell Markers, Stem Cells
Molecular Family	CD Molecules, Cytokine/Chemokine Receptors, GPCR
Antigen References	<ol style="list-style-type: none">1. Berger E, <i>et al.</i> 1999. <i>Annu. Rev. Immunol.</i> 17:657.2. Loetscher P, <i>et al.</i> 2000. <i>Adv. Immunol.</i> 74:127.3. Murphy P, <i>et al.</i> 2000. <i>Pharmacol. Rev.</i> 52:145.
Gene ID	7852

Related Protocols

[TotalSeq™-A Antibodies and Cell Hashing with 10x Single Cell 3' Reagent Kit v3 3.1 Protocol](#)

Other Formats

APC anti-human CD184 (CXCR4), Biotin anti-human CD184 (CXCR4), PE anti-human CD184 (CXCR4), PE/Cyanine5 anti-human CD184 (CXCR4), Purified anti-human CD184 (CXCR4), PerCP/Cyanine5.5 anti-human CD184 (CXCR4), PE/Cyanine7 anti-human CD184 (CXCR4), Brilliant Violet 421™ anti-human CD184 (CXCR4), Brilliant Violet 605™ anti-human CD184 (CXCR4), Purified anti-

human CD184 (CXCR4) (Maxpar® Ready), PE/Dazzle™ 594 anti-human CD184 (CXCR4), APC/Cyanine7 anti-human CD184 (CXCR4), Brilliant Violet 785™ anti-human CD184 (CXCR4), TotalSeq™-A0366 anti-human CD184 (CXCR4), TotalSeq™-C0366 anti-human CD184 (CXCR4), Brilliant Violet 510™ anti-human CD184 (CXCR4), Ultra-LEAF™ Purified anti-human CD184 (CXCR4), APC/Fire™ 750 anti-human CD184 (CXCR4), TotalSeq™-B0366 anti-human CD184 (CXCR4), TotalSeq™-D0366 anti-human CD184 (CXCR4)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587