

TotalSeq™-D0574 anti-human CD235a (Glycophorin A) Antibody

Catalog# / Size	349125 / 10 µg
Clone	HI264
Regulatory Status	RUO
Workshop	VII 70312
Other Names	PAS-2, Sialoglycoprotein alpha, MN sialoglycoprotein, Glycophorin A, MNS blood group
Isotype	Mouse IgG2a, κ
Barcode Sequence	AGAGTATGTATGGGA
Description	CD235a (Glycophorin A) is member of the glycophorin A family. It is a type I sialoglycoprotein with a molecular weight of 10 kD, present in the cell membrane as a homodimer. Glycophorin A is expressed by erythroid precursors and erythrocytes. It carries the antigen determinants for the MNS blood groups and has been proposed to be an inhibitor of hemagglutination and hemolysis. Glycophorin A binds siglec 5, the erythrocyte binding antigen (EBA-175) of <i>P. falciparum</i> and some viruses, including influenza virus and hepatitis A virus.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA
Preparation	The antibody was purified by chromatography and conjugated with TotalSeq™-D oligomer under optimal conditions.
Concentration	0.5 mg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.
Application	PG - Quality tested
Recommended Usage	<p>Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis and the oligomer sequence is confirmed by sequencing. TotalSeq™-D antibodies are compatible with Mission Bio's Tapestry Single-Cell Sequencing Platform for simultaneous detection of DNA and Protein.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
Additional Product Notes	<p>TotalSeq™-D reagents are designed to profile protein expression at single cell level. The Mission Bio Tapestry platform and sequencer (e.g. Illumina analyzers) are required. Please contact technical support for more information, or visit biolegend.com/totalseq/single-cell-dna</p> <p>The barcode flanking sequences are CGAGATGACTACGCTACTCATGG (PCR handle), and GAGCCGATCTAGTATCTCAGT*C*G (capture sequence). * indicates a phosphorothioated bond, to prevent nuclease degradation.</p>

View more applications data for this product in our [Application Technical Notes](#).

Application References

(PubMed link indicates BioLegend citation)

1. Mason D, *et al.* Eds. 2002. Leucocyte Typing VII: White Cell Differentiation Antigens. Oxford University Press. (FC)

RRID AB_2904370 (BioLegend Cat. No. 349125)

Antigen Details

Structure	Member of the glycoprotein A family; type I protein with a molecular weight of 10 kD; present in the membrane as homodimer
Distribution	Erythrocytes and erythroid precursors
Function	Possible inhibitor of hemagglutination and hemolysis; carries the antigen determinants for the MNS blood groups
Ligand/Receptor	Siglec 5, erythrocyte binding antigen (EBA-175) of <i>P. falciparum</i> , receptor for the influenza virus, hepatitis A virus
Cell Type	Erythrocytes
Biology Area	Cell Adhesion, Cell Biology, Immunology
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	<ol style="list-style-type: none">1. Reid ME. 2009. <i>Immunohematology</i> 25:95.2. Palacajornsuk P. 2006. <i>Immunohematology</i> 22:171.3. Pasvol G. 2003. <i>Trends Parasitol.</i> 19:430.4. Takakuwa Y. 2001. <i>Curr. Opin. Hematol.</i> 8:80.
Gene ID	2993

Related Protocols

[TotalSeq™-D with Mission Bio Tapestri® Single-Cell DNA + Protein Protocol](#)

Other Formats

Pacific Blue™ anti-human CD235a (Glycophorin A), PE anti-human CD235a (Glycophorin A), Purified anti-human CD235a (Glycophorin A), FITC anti-human CD235a (Glycophorin A), PerCP/Cyanine5.5 anti-human CD235a (Glycophorin A), PE/Cyanine7 anti-human CD235a (Glycophorin A), APC anti-human CD235a (Glycophorin A), APC/Cyanine7 anti-human CD235a (Glycophorin A), TotalSeq™-A0574 anti-human CD235a (Glycophorin A), PE/Dazzle™ 594 anti-human CD235a (Glycophorin A), TotalSeq™-C0574 anti-human CD235a (Glycophorin A), Alexa Fluor® 700 anti-human CD235a (Glycophorin A), TotalSeq™-D0574 anti-human CD235a (Glycophorin A)

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587