

True-Nuclear™ One Step Staining Mouse Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD25 PE/CD4 PerCP)

Catalog# / Size	136803 / 25 tests
Clone	MF-14
Regulatory Status	RUO
Other Names	Forkhead box protein P3, Scurfin, JM2, IPEX, Zinc finger protein JM2
Isotype	Rat IgG2b, κ
Description	FOXP3 is a 47 kD transcription factor, also known as Forkhead box protein P3, Scurfin, JM2, or IPEX. It is proposed to be a master regulatory gene and more specific marker of T regulatory cells than most cell surface markers (such as CD4 and CD25). Transduced expression of FOXP3 in CD4 ⁺ /CD25 ⁻ cells has been shown to induce GITR, CD103, and CTLA4 and impart a T regulatory cell phenotype. FOXP3 is mutated in X-linked autoimmunity-allergic dysregulation syndrome (XLAAD or IPEX) in humans and in "scurfy" mice. Overexpression of FOXP3 has been shown to lead to a hypoactive immune state suggesting that this transcriptional factor is a central regulator of T cell activity.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	This kit is guaranteed for six months. Upon receipt, store between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	<p>Materials Provided:</p> <ol style="list-style-type: none">1. Alexa Fluor® 488 anti-mouse FOXP3/CD25 PE/CD4 PerCP antibody cocktail - 25 tests2. Alexa Fluor® 488 Rat IgG2b, κ isotype control/CD25 PE/CD4 PerCP antibody cocktail - 25 tests3. True-Nuclear™ Transcription Buffer Set - 120 tests (Cat. No. 424401) <p>Materials not included:</p> <ol style="list-style-type: none">1. Cell Staining Buffer (Cat. No. 420201)2. Single Color Compensation Controls <p>Immunofluorescence Staining Procedures:</p> <ol style="list-style-type: none">1. Aliquot 100 µL of target cells to each tube.2. Add 1 mL of the Transcription Factor 1X Fix solution to each tube, vortex, and incubate at room temperature in the dark for 45-60 minutes.3. Without washing, add 2 mL of the Transcription Factor 1X Perm Buffer to each tube.4. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.5. Add 2 mL of the Transcription Factor 1X Perm Buffer to each tube.6. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.7. Resuspend the cell pellet in 100 µL of the Transcription Factor 1X Perm Buffer.8. Add 20 µL of Alexa Fluor® 488 anti-mouse FOXP3/CD25 PE/CD4 PerCP antibody cocktail or 20 µL of Alexa Fluor® 488 rat IgG2b, ? isotype control/CD25 PE/CD4 PerCP antibody cocktail into the appropriate tubes. Incubate in the dark at room temperature for at least 30 minutes.9. Add 2 mL of the Transcription Factor 1X Perm Buffer to each tube.10. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.11. Add 2 mL of the cell staining buffer.12. Centrifuge tubes at 400 x g at room temperature for five minutes, and discard the supernatant.13. Resuspend in 0.5 mL cell staining buffer and then acquire tubes on a flow cytometer.

Caution: The True-Nuclear™ Transcription Factor Buffer Set contains paraformaldehyde, which is

toxic and mutagenic. Please handle with caution. Wear gloves, lab coats, and necessary protection to avoid direct contact.

NOTE: For flow cytometric staining with this clone, True-Nuclear™ Transcription Factor Buffer Set (Cat. No. [424401](#)) offers improved staining and is highly recommended over the Foxp3/Perm Buffer Set (Cat. No. [421403](#)).

Application References

(PubMed link indicates BioLegend citation)

1. Ono M, *et al.* 2007. *Nature* 446:685.
2. Hori S, *et al.* 2003. *Science* 299:1057.
3. Fontenot JD, *et al.* 2003 *Nature Immunol* 4:330.
4. Fallarino F, *et al.* 2009. *J. Immunol.* 183:6033. [PubMed](#)
5. Barber A, *et al.* 2009 *J. Immunol.* 183:6939. [PubMed](#)
6. Nakashima H, *et al.* 2010. *J. Immunol.* 184:4637. [PubMed](#)

Product Citations

1. Peng K, *et al.* 2017. PLoS One.. 10.1371/journal.pone.0183271. [PubMed](#)
2. Lu Y, *et al.* 2018. *Cancer Cell.* 33:1048. [PubMed](#)


Antigen Details

Structure	50-55 kd protein. Forkhead/winged-helix transcription factor family, contains zinc finger and forkhead domains.
Distribution	Nuclear; expressed in Treg cells.
Function	Master regulatory gene in Treg cell development, crucial for immune homeostasis.
Interaction	Interacts with DNA
Cell Type	Tregs
Biology Area	Immunology, Transcription Factors
Molecular Family	CD Molecules, Nuclear Markers
Antigen References	<ol style="list-style-type: none">1. Ono M, <i>et al.</i> 2007. <i>Nature</i> 446:685.2. Hori S, <i>et al.</i> 2003. <i>Science</i> 299:1057.3. Fontenot JD, <i>et al.</i> 2003 <i>Nature Immunol</i> 4:330.
Regulation	Present at high level in T reg cells. Induced by T cell activation.
Gene ID	20371


Other Formats

Purified anti-mouse FOXP3, PE anti-mouse FOXP3, Alexa Fluor® 488 anti-mouse FOXP3, Alexa Fluor® 647 anti-mouse FOXP3, Pacific Blue™ anti-mouse FOXP3, Brilliant Violet 421™ anti-mouse FOXP3, True-Nuclear™ One Step Staining Mouse Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD25 PE/CD4 PerCP), Alexa Fluor® 700 anti-mouse FOXP3

Product Data



C57BL/6 splenocytes were stained with True-Nuclear™ One Step Staining Mouse Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD25 PE/CD4 PerCP).



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587