

Ultra-LEAF™ Purified anti-mouse IL-4 Antibody

Catalog# / Size	504121 / 100 µg 504122 / 1 mg 504135 / 5 mg 504136 / 25 mg 504137 / 50 mg 504138 / 100 mg
Clone	11B11
Regulatory Status	RUO
Other Names	Interleukin-4, Ia inducing factor (IaIF), B cell stimulating factor-1 (BSF-1), Hodgkin's cell growth factor (HCGF), Mast cell growth factor-2 (MCGF-2), Macrophage fusion factor (MFF), T cell growth factor-2 (TCGF-2)
Isotype	Rat IgG1, κ
Description	IL-4 is a pleiotropic cytokine produced by activated T cells, mast cells, and basophils. IL-4 is a potent lymphoid cell growth factor which stimulates the growth and activation of certain B cells and T cells. IL-4 is important for regulation of T helper subset development.

Product Details

Verified Reactivity	Mouse
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	Partially purified native mouse IL-4
Formulation	0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.01 EU/µg of the protein (<0.001 ng/µg of the protein) as determined by the LAL test.
Preparation	The Ultra-LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.
Concentration	The antibody is bottled at the concentration indicated on the vial, typically between 2 mg/mL and 3 mg/mL. Older lots may have also been bottled at 1 mg/mL. To obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C. This Ultra-LEAF™ solution contains no preservative; handle under aseptic conditions.
Application	ELISA Capture - Quality tested CyTOF® - Verified ELISPOT Capture, Neut, ICC, IP, IHC - Reported in the literature, not verified in house
Recommended Usage	Each lot of this antibody is quality control tested by ELISA assay . For ELISPOT applications, a concentration range of 2-6 µg/ml is recommended. For ELISA capture applications, a concentration range of 0.5-2.0 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of IL-4 recombinant protein ranging from 250 to 2 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes	ELISA^{1,2,10,13} or ELISPOT⁵ Capture: The purified 11B11 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated BVD6-24G2 antibody (Cat. No. 504202) as the detecting antibody and recombinant mouse IL-4 (Cat. No. 575609) as the standard. The LEAF™ purified antibody is suggested for ELISPOT capture. Neutralization^{1-2,9,12}: The 11B11 antibody can neutralize the bioactivity of natural or recombinant IL-4. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for neutralization of mouse IL-4 bioactivity <i>in vivo</i> and <i>in vitro</i> (Cat. No. 504108). Additional reported applications (for the relevant formats) include: immunoprecipitation ¹⁶ , immunohistochemical staining of formalin-fixed paraffin-embedded tissue sections ⁸ and paraformaldehyde-fixed, saponin-treated frozen tissue sections ^{6,7} , and immunocytochemistry ⁴ . Note: For testing mouse IL-4 in serum, plasma or supernatant, BioLegend's ELISA Max™ Sets

(Cat. No. 431101 to 431106) are specially developed and recommended.

Additional Product Notes

Get a 50% discount on this product when purchased in our Activation Bundles. Restrictions apply. [Learn more...](#)

Application References

(PubMed link indicates BioLegend citation)

1. Shirai A, *et al.* 1994. *Cytokine* 6:329. (ELISA, Neut)
2. Abrams J. 1995. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.20. (ELISA, Neut)
3. Assenmacher M, *et al.* 1994. *Eur. J. Immunol.* 24:1097.
4. Openshaw P, *et al.* 1995. *J. Exp. Med.* 182:1357. (ICC)
5. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.19. (ELISA Capture)
6. Litton M, *et al.* 1994. *J. Immunol. Methods* 175:47. (IHC)
7. Andersson U, *et al.* 1999. *Detection and quantification of gene expression.* New York:Springer-Verlag. (IHC)
8. Fan WY, *et al.* 2001. *Exp. Biol. Med.* 226:1045. (IHC)
9. Hara M, *et al.* 2001. *J. Immunol.* 166:3789. (Neut)
10. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. (ELISA)
11. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366.
12. Wang W, *et al.* 2007. *J. Immunol.* 178:4885. (Neut)
13. Xu G, *et al.* 2007. *J. Immunol.* 179:5358. (ELISA) [PubMed](#)
14. Ohnmacht C, *et al.* 2008. *Blood* 113:2816. [PubMed](#)
15. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
16. Zavorotinskaya T, *et al.* 2003. *Mol. Ther.* 7:155. (IP)

Product Citations

1. Ogbечи J, *et al.* 2022. *Front Immunol.* 13:1001956. [PubMed](#)
2. Kenichi Shimada *et al.* 2018. *Immunity.* 49(5):873-885. [PubMed](#)
3. Glaubitz J, *et al.* 2022. *Nat Commun.* 13:4502. [PubMed](#)
4. Imanishi T, *et al.* 2020. *Cell Reports.* 32(3):107911. [PubMed](#)
5. Jia X, *et al.* 2021. *Front Cell Dev Biol.* 9:685522. [PubMed](#)
6. Montes de Oca M, *et al.* 2020. *PLoS Pathog.* 16:e1008994. [PubMed](#)
7. Guo P, *et al.* 2021. *J Immunol.* 207:408. [PubMed](#)
8. Kanno T, *et al.* 2022. *Front Immunol.* 13:904875. [PubMed](#)
9. Xia X, *et al.* 2020. *J Clin Invest.* 130:5180. [PubMed](#)
10. Hao L, *et al.* 2021. *Br J Pharmacol.* 178:4726. [PubMed](#)
11. Price C, *et al.* 2020. *Cell Host & Microbe.* 27(4):571-584.e7. [PubMed](#)
12. Okubo A, *et al.* 2021. *Int J Mol Sci.* 23: [PubMed](#)
13. Ding X, *et al.* 2021. *Front Immunol.* 12:667136. [PubMed](#)

RRID

AB_11149679 (BioLegend Cat. No. 504121)
AB_11150601 (BioLegend Cat. No. 504122)
AB_2750404 (BioLegend Cat. No. 504135)
AB_2750405 (BioLegend Cat. No. 504136)
AB_2750406 (BioLegend Cat. No. 504137)
AB_2750407 (BioLegend Cat. No. 504138)

Antigen Details

Structure	Cytokine; 15-19 kD (Mammalian)
Bioactivity	Differentiation of naïve CD4 ⁺ T cells to the T _H 2 type, proliferation/differentiation of activated B cells, expression of class II MHC antigens, and of low affinity IgE receptors in resting B cells
Cell Sources	Mast cells, T cells, bone marrow stromal cells
Cell Targets	B cells, T cells, monocytes, endothelial cells, fibroblasts
Receptors	Heterodimer IL-4R α (CD124); γ -subunit (CD132) in common with IL-2R, IL-7R, IL-13R, IL-15R
Cell Type	Tregs
Biology Area	Immunology
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press San Diego.2. Boulay J, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:294.3. Dullens H, <i>et al.</i> 1991. <i>In vivo</i> 5:567.4. Paul W. 1991. <i>Blood</i> 77:1859.
Regulation	Upregulated by IL-2, platelet activating factor; downregulated by TGF- β
Gene ID	16189

Related Protocols


Active Protocols: [Sandwich ELISA - Video](#)

[Sandwich ELISA Protocol](#)

Other Formats

APC anti-mouse IL-4, PE anti-mouse IL-4, Purified anti-mouse IL-4, Alexa Fluor® 488 anti-mouse IL-4, Alexa Fluor® 647 anti-mouse IL-4, PE/Cyanine7 anti-mouse IL-4, Brilliant Violet 421™ anti-mouse IL-4, Ultra-LEAF™ Purified anti-mouse IL-4, PerCP/Cyanine5.5 anti-mouse IL-4, Brilliant Violet 605™ anti-mouse IL-4, Purified anti-mouse IL-4 (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse IL-4, Brilliant Violet 711™ anti-mouse IL-4, APC/Fire™ 750 anti-mouse IL-4

Product Data



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587