

## Alexa Fluor® 647 anti-mouse IL-10 Antibody

<b>Catalog# / Size</b>	505016 / 25 µg 505014 / 100 µg
<b>Clone</b>	JES5-16E3
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Interleukin-10, Cytokine synthesis inhibitory factor (CSIF), B cell derived T cell growth factor (B-TCGF)
<b>Isotype</b>	Rat IgG2b, κ
<b>Description</b>	IL-10 was originally described as Cytokine Synthesis Inhibitory Factor (CSIF) by virtue of its ability to inhibit cytokine production by Th1 clones. IL-10 shares over 80% sequence homology with the Epstein-Barr virus protein BCRF1. IL-10 inhibits IFN-γ, TNF-β, and IL-2 production by Th1 clones; inhibits macrophage-mediated IL-1, IL-6, and TNF-α synthesis; suppresses the delayed type hypersensitivity response; stimulates Th2 cell response (which results in elevated antibody production); and promotes mast cell proliferation in combination with IL-4.

### Product Details

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	<i>E. coli</i> -expressed, recombinant mouse IL-10
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">ICFC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">intracellular immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is ≤ 0.25 µg per 10 <sup>6</sup> cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.  * Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.  Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.  <a href="#">View full statement regarding label licenses</a>
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Application Notes</b>	<b>ELISA or ELISPOT Detection</b> <sup>1,9,11</sup> : The biotinylated JES5-16E3 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified JES5-2A5 antibody (Cat. Nos. 504902 & 504904) as the capture antibody. <b>ELISA Capture</b> : The purified JES5-16E3 antibody is useful as the capture antibody in a sandwich ELISA when used in conjunction with the biotinylated JES5-2A5 antibody (Cat. No. 505003) as the detection antibody and recombinant mouse IL-10 (Cat. No. 575809) as the standard. <b>Neutralization</b> <sup>14</sup> : The Ultra-LEAF™ purified JES5-16E3 antibody can neutralize the bioactivity of natural or recombinant IL-10. <b>Flow Cytometry</b> <sup>3</sup> : The fluorochrome-labeled JES5-16E3 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-10-producing cells within mixed cell populations. <b>Additional reported applications (for relevant formats) include</b> : immunohistochemistry <sup>3</sup> .

## Application References

1. Simkin G, *et al.* 2000. *J. Immunol.* 164:2457.
2. Kitagaki K, *et al.* 2002. *Clin. Diagn. Lab Immunol.* 9:1260.
3. Khanna A, *et al.* 2000. *J. Immunol.* 164:1346.
4. Sander B, *et al.* 1993. *J. Immunol. Methods* 166:201.
5. Litton M, *et al.* 1994. *J. Immunol. Methods* 175:47.
6. Andersson U, *et al.* 1999. *Detection and quantification of gene expression.* New York:Springer-Verlag.
7. Finkelman F, *et al.* 2003. *Curr. Prot. Immunol.* John Wiley & Sons New York. Unit 6.28.
8. Wang W, *et al.* 2004. *FASEB J.* 18:1043.
9. Brummel R and Lenert P. 2005. *J. Immunol.* 174:2429.
10. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366.
11. Xu G, *et al.* 2007. *J. Immunol.* 179:5358. [PubMed](#)
12. Brummel R, *et al.* 2005. *J. Immunol.* 174:2429. [PubMed](#)
13. Kang YJ, *et al.* 2007. *Stem Cells* 25:1814. [PubMed](#)
14. Seo N, *et al.* 2001. *Immunology.* 103:449. (Neut)

## Product Citations

1. Wu J *et al.* 2017. *Immunity.* 47(6):1114-1128. [PubMed](#)
2. Tuganbaev T, *et al.* 2020. *Cell.* 182(6):1441-1459.e21. [PubMed](#)
3. Jain RW, *et al.* 2018. *Cell Rep.* 25:3342. [PubMed](#)
4. Rio R, *et al.* 2012. *J Immunol.* 188:541. [PubMed](#)
5. Nasti A, *et al.* 2017. *Eur J Immunol.* 47:2163. [PubMed](#)

## RRID

AB\_2125096 (BioLegend Cat. No. 505016)  
AB\_493511 (BioLegend Cat. No. 505014)

## Antigen Details

---

<b>Structure</b>	Acid-labile cytokine, dimer, 17-21 kD (Mammalian)
<b>Cell Sources</b>	Activated CD8 <sup>+</sup> T cells, Th0, Th2 subset of CD4 <sup>+</sup> T cells, Ly-1 <sup>+</sup> B cells, monocytes, macrophages, keratinocytes
<b>Cell Targets</b>	T cells, B cells, mast cells, macrophages
<b>Receptors</b>	IL-10R (CDw210)
<b>Cell Type</b>	Tregs
<b>Biology Area</b>	Immunology
<b>Molecular Family</b>	Cytokines/Chemokines
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press San Diego.</li><li>2. de Waal-Malefy R, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:314.</li><li>3. Howard M, <i>et al.</i> 1992. <i>Immunol. Today</i> 13:198.</li><li>4. Quesniaux V. 1992. <i>Res. Immunol.</i> 143:385.</li><li>5. Norton SK, <i>et al.</i> 2008. <i>J. Immunol.</i> 180:2848.</li></ol>
<b>Regulation</b>	Downregulated by IL-4, IL-10
<b>Gene ID</b>	<a href="#">16153</a>

## Related Protocols

---

[Surface and Intracellular Cytokine Staining for Flow Cytometry - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)


## Other Formats

---

APC anti-mouse IL-10, Biotin anti-mouse IL-10, FITC anti-mouse IL-10, PE anti-mouse IL-10, Purified anti-mouse IL-10, Alexa Fluor® 647 anti-mouse IL-10, PE/Cyanine7 anti-mouse IL-10, Alexa Fluor® 488 anti-mouse IL-10, Brilliant Violet 421™ anti-mouse IL-10, Pacific Blue™ anti-mouse IL-10, PerCP/Cyanine5.5 anti-mouse IL-10, Purified anti-mouse IL-10 (Maxpar® Ready), Brilliant Violet 605™ anti-mouse IL-10, PE/Dazzle™ 594 anti-mouse IL-10, APC/Cyanine7 anti-mouse IL-10, Ultra-LEAF™ Purified anti-mouse IL-10

## Product Data

---



PMA/ionomycin-stimulated Th2-polarized Balb/c mouse splenocytes were intracellularly stained with JES5-16E3 Alexa Fluor® 647 and CD3 (17A2) PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587