

APC anti-human CD14 Antibody

| | |
|--------------------------|---|
| Catalog# / Size | 301807 / 25 tests 301808 / 100 tests |
| Clone | M5E2 |
| Regulatory Status | RUO |
| Workshop | III 329 |
| Other Names | LPS receptor |
| Isotype | Mouse IgG2a, κ |
| Description | CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein also known as LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens, and in the upregulation of adhesion molecules and expression of cytokines in monocytes and neutrophils. |

Product Details

| | |
|-------------------------------|--|
| Verified Reactivity | Human, Cynomolgus, Rhesus |
| Reported Reactivity | African Green, Capuchin Monkey, Cow, Chimpanzee, Common Marmoset, Cotton-topped Tamarin, Dog, Pigtailed Macaque, Squirrel Monkey |
| Antibody Type | Monoclonal |
| Host Species | Mouse |
| Immunogen | Full-length human CD14 protein |
| Formulation | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA) |
| Preparation | The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. |
| Concentration | Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.) |
| Storage & Handling | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze. |
| Application | FC - Quality tested |
| Recommended Usage | Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. |
| Excitation Laser | Red Laser (633 nm) |
| Application Notes | The M5E2 antibody inhibits monocyte activation and cytokine production induced by LPS. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections, blocking of LPS stimulation ⁴ , and immunofluorescence microscopy ⁵ . Clone M5E2 is not recommended for immunohistochemical staining of formalin-fixed paraffin-embedded sections. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 301861 and 301862). |
| Application References | <ol style="list-style-type: none"> 1. McMichael A, <i>et al.</i> 1987. Leucocyte Typing III. Oxford University Press. New York. 2. Knapp W, <i>et al.</i> Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York. (IHC-F) 3. Schlossman S, <i>et al.</i> Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. 4. Power CP, <i>et al.</i> 2004. <i>J. Immunol.</i> 173:5229. (Block) 5. Williams KC, <i>et al.</i> 2001. <i>J. Exp. Med.</i> 193:905. |

6. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (FC) [PubMed](#)
7. Santer DM, *et al.* 2010. *J. Immunol.* 485:4739. [PubMed](#)
8. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
9. Zizzo G, *et al.* 2012. *J. Immunol.* 189:3508. [PubMed](#)
10. Stoeckius M, *et al.* 2017. *Nat. Methods.* 14:865. (PG)
11. Peterson VM, *et al.* 2017. *Nat. Biotechnol.* 35:936. (PG)

Product Citations

1. Zhu ZY, *et al.* 2022. *Front Immunol.* 12:781087. [PubMed](#)
2. Mahmud FJ, *et al.* 2020. *J Neuroinflammation.* 0.897916667. [PubMed](#)
3. Lopes MG, *et al.* 2021. *J Dairy Sci.* 104:9340. [PubMed](#)
4. Woolsey C, *et al.* 2020. *Scientific Reports.* 10(1):3071.. [PubMed](#)
5. Alharthi AS, *et al.* 2019. *J Dairy Sci.* 102:10599. [PubMed](#)
6. Downes DJ, *et al.* 2021. *Nat Genet.* 53:1606. [PubMed](#)
7. Offersen R, *et al.* 2016. *J Virol.* 90: 4441 - 4453. [PubMed](#)
8. Woolsey C *et al.* 2019. *Cell Rep.* 28(12):3032-3046 . [PubMed](#)
9. Kedage V, *et al.* 2019. *MAbs.* 12:1685832. [PubMed](#)
10. Guo Z, *et al.* 2022. *NPJ Precis Oncol.* 6:34. [PubMed](#)
11. Ferrari M, *et al.* 2021. *J Virol.* 95:e0068521. [PubMed](#)
12. Jankeel A, *et al.* 2020. *J Virol.* :94. [PubMed](#)
13. Constantinescu-Bercu A, *et al.* 2020. *eLife.* 9:00. [PubMed](#)
14. Yokoyama K, *et al.* 2022. *Front Immunol.* 13:1016914. [PubMed](#)
15. Spildrejorde M, *et al.* 2014. *Physiol Genomics.* 46:512. [PubMed](#)
16. Mu Z, *et al.* 2021. *Genome Biol.* 122:22. [PubMed](#)
17. Woodberry T, *et al.* 2017. *Infect Immun.* 85:e00986. [PubMed](#)
18. Takele Y, *et al.* 2022. *PLoS Negl Trop Dis.* 16:e0010544. [PubMed](#)
19. Broeren MGA, *et al.* 2018. *ALTEX.* 36:18:00. [PubMed](#)
20. Claus C, *et al.* 2021. *Pharmaceutics.* 13:. [PubMed](#)
21. Wegner J, *et al.* 2021. *Sci Rep.* 11:14983. [PubMed](#)
22. da Rocha Sobrinho HM, *et al.* 2021. *Journal of Neuroscience Research.* .: [PubMed](#)
23. Powell EJ, *et al.* 2018. *J Dairy Sci.* 101:8182. [PubMed](#)
24. Ferguson Bennit HR, *et al.* 2017. *Blood Lymphat Cancer.* 7:9. [PubMed](#)
25. Zhou Z, *et al.* 2018. *J Dairy Sci.* 101:10374. [PubMed](#)
26. Weiss R, *et al.* 2021. *Cells.* 10:. [PubMed](#)
27. Chen Y, *et al.* 2021. *Front Mol Biosci.* 8:777370. [PubMed](#)
28. Kiers D *et al.* 2018. *EBioMedicine.* 33:144-156 . [PubMed](#)
29. Lavaert M, *et al.* 2020. *Immunity.* 52(6):1088-1104. [PubMed](#)
30. Tähtinen S, *et al.* 2022. *Nat Immunol.* 23:532. [PubMed](#)
31. Bade RM, *et al.* 2021. *Molecular Oncology.* . [PubMed](#)
32. Miksiunas R, *et al.* 2020. *Int J Mol Sci.* 21:00. [PubMed](#)
33. Duncan CJA, *et al.* 2022. *J Exp Med.* 219:. [PubMed](#)
34. Wang H, *et al.* 2017. *J Neurochem.* 141:347. [PubMed](#)
35. Alhaj Hussien K, *et al.* 2020. *Front Immunol.* 11:579776. [PubMed](#)
36. Thomas BL, *et al.* 2022. *J Dairy Sci.* 105:2256. [PubMed](#)
37. Magg T, *et al.* 2021. *Sci Immunol.* 6:. [PubMed](#)

RRID

AB_314189 (BioLegend Cat. No. 301807)
 AB_314190 (BioLegend Cat. No. 301808)

Antigen Details


| | |
|---------------------------|---|
| Structure | GPI-linked membrane glycoprotein, 53-55 kD |
| Distribution | Monocytes, macrophages, granulocytes (low) |
| Function | LPS receptor, clearance of Gram-negative pathogens |
| Ligand/Receptor | LPS |
| Cell Type | Granulocytes, Macrophages, Monocytes, Neutrophils |
| Biology Area | Cell Biology, Immunology, Innate Immunity, Neuroinflammation, Neuroscience |
| Molecular Family | CD Molecules |
| Antigen References | <ol style="list-style-type: none"> 1. Stocks S, <i>et al.</i> 1990. <i>Biochem. J.</i> 268:275. 2. Wright S, <i>et al.</i> 1990. <i>Science</i> 249:1434. |
| Gene ID | 929 |

Related Protocols

Other Formats

APC anti-human CD14, FITC anti-human CD14, PE anti-human CD14, Purified anti-human CD14, PE/Cyanine7 anti-human CD14, Alexa Fluor® 488 anti-human CD14, Alexa Fluor® 647 anti-human CD14, Ultra-LEAF™ Purified anti-human CD14, Pacific Blue™ anti-human CD14, APC/Cyanine7 anti-human CD14, Alexa Fluor® 700 anti-human CD14, PerCP/Cyanine5.5 anti-human CD14, Biotin anti-human CD14, Brilliant Violet 421™ anti-human CD14, Brilliant Violet 570™ anti-human CD14, Brilliant Violet 605™ anti-human CD14, Brilliant Violet 650™ anti-human CD14, Brilliant Violet 711™ anti-human CD14, Brilliant Violet 785™ anti-human CD14, Brilliant Violet 510™ anti-human CD14, Purified anti-human CD14 (Maxpar® Ready), PerCP anti-human CD14, PE/Dazzle™ 594 anti-human CD14, APC/Fire™ 750 anti-human CD14, TotalSeq™-A0081 anti-human CD14, TotalSeq™-B0081 anti-human CD14, TotalSeq™-C0081 anti-human CD14, PE/Cyanine5 anti-human CD14, TotalSeq™-D0081 anti-human CD14, GMP FITC anti-human CD14, GMP APC anti-human CD14, GMP PE anti-human CD14, GMP Pacific Blue™ anti-human CD14

Product Data



Human peripheral blood monocytes stained with M5E2 APC

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587