

PE/Cyanine7 anti-human TCR V δ 2 Antibody

Catalog# / Size	331421 / 25 tests 331422 / 100 tests
Clone	B6
Regulatory Status	RUO
Other Names	T cell receptor V δ 2
Isotype	Mouse IgG1, κ
Description	The V δ 2 TCR is a variant of the TCR δ chain expressed on a subset of γ/δ T cells. V γ 9V δ 2 T lymphocytes, a major γ/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomorphic proteins CD3 δ , γ , ϵ , and ζ chains.

Product Details

Verified Reactivity	Human
Reported Reactivity	Chimpanzee, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Cyanine7 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Additional Product Notes	BioLegend is in the process of converting the name PE/Cy7 to PE/Cyanine7. The dye molecule remains the same, so you should expect the same quality and performance from our PE/Cyanine7 products. Please contact Technical Service if you have any questions.
Application References	<ol style="list-style-type: none">1. Rojas RE, <i>et al.</i> 2005. <i>J. Infect. Dis.</i> 192:1806.2. Correia DV, <i>et al.</i> 2011. <i>Blood</i> 118:992. (FC) PubMed
Product Citations	<ol style="list-style-type: none">1. Cano CE, <i>et al.</i> 2021. <i>Cell Reports.</i> 36(2):109359. PubMed
RRID	AB_2687327 (BioLegend Cat. No. 331421) AB_2687328 (BioLegend Cat. No. 331422)

Antigen Details

Structure	Ig superfamily member, TCR γ/δ with CD3 forms the CD3/TCR complex
Distribution	Expressed on majority of peripheral γ/δ T cells
Function	Antigen recognition, T cell activation
Ligand/Receptor	Peptide bound to MHC
Cell Type	T cells
Biology Area	Adaptive Immunity, Immunology
Molecular Family	TCRs
Antigen References	1. Scotet E, <i>et al.</i> 2005. <i>Immunity</i> 22:71. 2. Rincon-Orozco B, <i>et al.</i> 2005. <i>J. Immunol.</i> 175:2144.
Gene ID	6964



Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human TCR V δ 2, Biotin anti-human TCR V δ 2, FITC anti-human TCR V δ 2, PE anti-human TCR V δ 2, PerCP anti-human TCR V δ 2, Brilliant Violet 711™ anti-human TCR V δ 2, Pacific Blue™ anti-human TCR V δ 2, Alexa Fluor® 700 anti-human TCR V δ 2, PerCP/Cyanine5.5 anti-human TCR V δ 2, APC anti-human TCR V δ 2, PE/Cyanine7 anti-human TCR V δ 2, APC/Fire™ 750 anti-human TCR V δ 2, Brilliant Violet 421™ anti-human TCR V δ 2, Brilliant Violet 605™ anti-human TCR V δ 2, PE/Dazzle™ 594 anti-human TCR V δ 2, Brilliant Violet 510™ anti-human TCR V δ 2, TotalSeq™-A0582 anti-human TCR V δ 2, TotalSeq™-C0582 anti-human TCR V δ 2, TotalSeq™-B0582 anti-human TCR V δ 2, APC/Cyanine7 anti-human TCR V δ 2

Product Data



Human peripheral blood lymphocytes were stained with CD3 APC and TCR V δ 2 (clone B6) PE/Cyanine7 (top) or mouse IgG1, κ isotype control (bottom).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all

applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587